

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



Spoon Feeding Reduction Methods



Simplified Knowledge Management Classes Bangalore

My name is [Subhashish Chattopadhyay](#). I have been teaching for IIT-JEE, Various International Exams (such as IMO [International Mathematics Olympiad], IPhO [International Physics Olympiad], IChO [International Chemistry Olympiad]), IGCSE (IB), CBSE, I.Sc, Indian State Board exams such as WB-Board, Karnataka PU-II etc since 1989. As I write this book in 2016, it is my 25 th year of teaching. I was a Visiting Professor to BARC Mankhurd, Chembur, Mumbai, Homi Bhabha Centre for Science Education (HBCSE) Physics Olympics camp BARC Campus.

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

I am Life Member of ...

- [IAPT \(Indian Association of Physics Teachers \)](#)
- [IPA \(Indian Physics Association \)](#)
- [AMTI \(Association of Mathematics Teachers of India \)](#)
- [National Human Rights Association](#)
- [Men's Rights Movement \(India and International \)](#)
- [MGTOW Movement \(India and International \)](#)

And also of

[IACT \(Indian Association of Chemistry Teachers \)](#)



The selection for National Camp (for Official Science Olympiads - Physics, Chemistry, Biology, Astronomy) happens in the following steps

1) **NSEP** (National Standard Exam in Physics) and **NSEC** (National Standard Exam in Chemistry) held around 24 rth November. Approx 35,000 students appear for these exams every year. The exam fees is Rs 100 each. Since 1998 the IIT JEE toppers have been topping these exams and they get to know their rank / performance ahead of others.

2) **INPhO** (Indian National Physics Olympiad) and **INChO** (Indian National Chemistry Olympiad). Around 300 students in each subject are allowed to take these exams. Students coming from outside cities are paid fair from the Govt of India.

3) The Top 35 students of each subject are invited at HBCSE (Homi Bhabha Center for Science Education) Mankhurd, near Chembur, BARC, Mumbai. After a 2-3 weeks camp the top 5 are selected to represent India. The flight tickets and many other expenses are taken care by Govt of India.

Since last 50 years there has been no dearth of “Good Books“. Those who are interested in studies have been always doing well. This e-Book does not intend to replace any standard text book. These topics are very old and already standardized.

There are 3 kinds of Text Books

- The thin Books - Good students who want more details are not happy with these. Average students who need more examples are not happy with these. Most students who want to “Cram” quickly and pass somehow find the thin books “good” as they have to read less !!

- The Thick Books - Most students do not like these, as they want to read as less as possible. Average students are “busy” with many other things and have no time to read all these.

- The Average sized Books - Good students do not get all details in any one book. Most bad students do not want to read books of “this much thickness“ also !!

We know there can be no shoe that’s fits in all.



Printed books are not e-Books! Can’t be downloaded and kept in hard-disc for reading “later”

So if you read this book later, you will get all kinds of examples in a single place. This becomes a very good “Reference Material”. I sincerely wish that all find this “very useful”.

Students who do not practice lots of problems, do not do well. The rules of “doing well” had never changed Will never change !

In this e-Book I am trying to solve this problem. Those students who practice can learn.

No one can help those who are not studying, or practicing.



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams


Learn more at <http://skmclasses.weebly.com/iit-jee-home-tuitions-bangalore.html>

Twitter - <https://twitter.com/ZookeeperPhy>

Facebook - <https://www.facebook.com/IIT.JEE.by.Prof.Subhashish/>

Blog - <http://skmclasses.kinja.com>

Blog - <http://skmclasses.blog.com>



A very polite request :

I wish these e-Books are read only by Boys and Men. Girls and Women, better read something else; learn from somewhere else.

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

Preface

We all know that in the species “Homo Sapiens “, males are bigger than females. The reasons are explained in standard 10, or 11 (high school) Biology texts. **This shapes or size, influences all of our culture.** Before we recall / understand the reasons once again, let us see some random examples of the influence

Random - 1

If there is a Road rage, then who all fight ? (generally ?). Imagine two cars driven by adult drivers. Each car has a woman of similar age as that of the Man. The cars “ touch “ or “ some issue happens”. Who all comes out and fights ? Who all are most probable to drive the cars ?



(Men are eager to fight, eager to rule, eager for war. Men want to drive. Men want to win)

Random - 2

Heavy metal music artists are all Men. Metallica, Black Sabbath, Motley Crue, Megadeth, Motorhead, AC/DC, Deep Purple, Slayer, Guns & Roses, Led Zeppelin, Aerosmith the list can be in thousands. All these are grown-up Boys, known as Men.



(Men strive for perfection. Men are eager to excel. Men work hard. Men want to win.)



Random - 3

Apart from Marie Curie, only one more woman got Nobel Prize in Physics. (Maria Goeppert Mayer - 1963). So, ... almost all are men.

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



(Men want to excel. Men strive for perfection. Men want to win. Men work hard. Men do better than women.)

Random - 4

The best Tabla Players are all Men.



(Men want to excel. Men strive for perfection. Men want to win. Men work hard. Men do better than women.)

Random - 5

History is all about, which Kings ruled. Kings, their men, and Soldiers went for wars. History is all about wars, fights, and killings by men.



Boys start fighting from school days. Girls do not fight like this

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



(Men are eager to fight, eager to rule, eager for war. Men want to drive. Men want to win.)

Random - 6

The highest award in Mathematics, the “ Fields Medal “ is around since decades. Till date only one woman could get that. (Maryam Mirzakhani - 2014). So, ... almost all are men.



(Men want to excel. Men strive for perfection. Men want to win. Men work hard. Men do better than women.)

Random - 7

Actor is a gender neutral word. Could the movie like “ Top Gun “ be made with Female actors ? The best pilots, astronauts, Fighters are all Men.



Random - 8

In my childhood had seen a movie named “ The Tower in Inferno “. In the movie when the tall tower is in fire, women were being saved first, as only one lift was working....



Many decades later another movie is made. A box office hit. “ The Titanic “. In this also As the ship is sinking women are being saved. **Men are disposable.** Men may get their turn later...



Movies are not training programs. Movies do not teach people what to do, or not to do. Movies only reflect the prevalent culture. Men are disposable, is the culture in the society. Knowingly, unknowingly, the culture is depicted in Movies, Theaters, Stories, Poems, Rituals, etc. I or you can't write a story, or make a movie in which after a minor car accident the Male passengers keep seating in the back seat, while the both the women drivers come out of the car and start fighting very bitterly on the road. There has been no story in this world, or no movie made, where after an accident or calamity, Men are being helped for safety first, and women are told to wait.

Random - 9

Artists generally follow the prevalent culture of the Society. In paintings, sculptures, stories, poems, movies, cartoon, Caricatures, knowingly / unknowingly, " the prevalent Reality " is depicted. The opposite will not go well with people. If deliberately " the opposite " is shown then it may only become a special art, considered as a special mockery.

पत्नी (सल्टू से): मुझे
नई साड़ी ला दो प्लीज।
सल्टू: पर तुम्हारी
दो-दो अलमारियां सा
डियों से ही तो भरी है।
पत्नी - वह सारी तो
पूरे मोहल्ले वालों ने
देख रखी है।
सल्टू - तो साड़ी लेने
के बजाए मोहल्ला
बदल लेते हैं।



Random - 10

Men go to "girl / woman's house" to marry / win, and bring her to his home. That is a sort of winning her. When a boy gets a " Girl-Friend ", generally he and his friends consider that as an achievement. The boy who " got / won " a girl-friend feels proud. His male friends feel, jealous, competitive and envious. Millions of stories have been written on these themes. Lakhs of movies show this. Boys / Men go for " bike race ", or say " Car Race ", where the winner " gets " the most beautiful girl of the college.



(Men want to excel. Men are eager to fight, eager to rule, eager for war. Men want to drive. Men want to win.)

Prithviraj Chauhan ' went ` to " pickup " or " abduct " or " win " or " bring " his love. There was a Hindi movie (hit) song ... " Pasand ho jaye, to ghar se utha laye ". It is not other way round. Girls do not go to Boy's house or man's house to marry. Nor the girls go in a gang to " pick-up " the boy / man and bring him to their home / place / den.

Random - 11

Rich people; often are very hard working. Successful business men, establish their business (empire), amass lot of wealth, with lot of difficulty. Lots of sacrifice, lots of hard work, gets into this. Rich people's wives had no contribution in this wealth creation. Women are smart, and successful upto the extent to choose the right/rich man to marry. So generally what happens in case of Divorces ? Search the net on " most costly divorces " and you will know. The women;(who had no contribution at all, in setting up the business / empire), often gets in Billions, or several Millions in divorce settlements.

Number 1

Rupert & Anna Murdoch -- \$1.7 billion

One of the richest men in the world, **Rupert Murdoch** developed his worldwide media empire when he inherited his father's Australian newspaper in 1952. He married Anna Murdoch in the '60s and they remained together for 32 years, springing off three children.

They split amicably in 1998 but soon Rupert forced Anna off the board of News Corp and the gloves came off. The divorce was finalized in June 1999 when Rupert agreed to let his ex-wife leave with \$1.7 billion worth of his assets, \$110 million of it in cash. Seventeen days later, Rupert married Wendi Deng, one of his employees.



Ted Danson & Casey Coates -- \$30 million

Ted Danson's claim to fame is undoubtedly his decade-long stint as Sam Malone on NBC's celebrated sitcom Cheers. While he did other TV shows and movies, he will always be known as the bartender of that place where everybody knows your name. He met his future first bride Casey, a designer, in 1976 while doing Erhard Seminars Training.

Ten years his senior, she suffered a paralyzing stroke while giving birth to their first child in 1979. In order to nurse her back to health, Danson took a break from acting for six months. But after two children and 15 years of marriage, the infatuation fell to pieces. Danson had started seeing Whoopi Goldberg while filming the comedy, Made in America and this precipitated the 1992 divorce. Casey got \$30 million for her trouble.

See <https://zookeepersblog.wordpress.com/misandry-and-men-issues-a-short-summary-at-single-place/>

See <http://skmclasses.kinja.com/save-the-male-1761788732>


It was Boys and Men, who brought the girls / women home. The Laws are biased, completely favoring women. The men are paying for their own mistakes.

See <https://zookeepersblog.wordpress.com/biased-laws/>

(Man brings the Woman home. When she leaves, takes away her share of big fortune!)

Random - 12

A standardized test of Intelligence will never be possible. It never happened before, nor ever will happen in future; where the IQ test results will be acceptable by all. In the net there are thousands of charts which show that the intelligence scores of girls / women are lesser. Debates of Trillion words, does not improve performance of Girls.



I am not wasting a single second debating or discussing with anyone, on this. I am simply accepting ALL the results. IQ is only one of the variables which is required for success in life. Thousands of books have been written on “ Networking Skills “, EQ (Emotional Quotient), Drive, Dedication, Focus, “ Tenacity towards the end goal “ ... etc. In each criteria, and in all together, women (in general) do far worse than men. Bangalore is known as “ capital of India “. [Fill in the blanks]. The blanks are generally filled as “ Software Capital “, “ IT Capital “, “ Startup Capital “, etc. I am member in several startup eco-systems / groups. I have attended hundreds of meetings, regarding “ technology startups “, or “ idea startups “. These meetings have very few women. Starting up new companies are all “ Men’s Game “ / “ Men’s business “. Only in Divorce settlements women will take their goodies, due to Biased laws. There is no dedication, towards wealth creation, by women.

Random - 13

Many men, as fathers, very unfortunately treat their daughters as “ Princess “. Every “ non-performing “ woman / wife was “ princess daughter “ of some loving father. Pampering the girls, in name of “ equal opportunity “, or “ women empowerment “, have led to nothing.



See <http://skmclasses.kinja.com/progressively-daughters-become-monsters-1764484338>

See <http://skmclasses.kinja.com/vivacious-vixens-1764483974>

There can be thousands of more such random examples, where “ Bigger Shape / size “ of males have influenced our culture, our Society. **Let us recall the reasons**, that we already learned in standard 10 - 11, Biology text Books. In humans, women have a long gestation period, and also spends many years (almost a decade) to grow, nourish, and stabilize the child. (Million years of habit) Due to survival instinct Males want to inseminate. Boys and Men fight for the “ facility (of womb + care) “ the girl / woman may provide. Bigger size for males, has a winning advantage. Whoever wins, gets the “ woman / facility “. The male who is of “ Bigger Size “, has an advantage to win.... Leading to Natural selection over millions of years. In general “ Bigger Males “; the “ fighting instinct “ in men; have led to wars,

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

and solving tough problems (Mathematics, Physics, Technology, startups of new businesses, Wealth creation, Unreasonable attempts to make things [such as planes], Hard work)

So let us see the IIT-JEE results of girls. Statistics of several years show that there are around 17, (or less than 20) girls in top 1000 ranks, at all India level. Some people will yet not understand the performance, till it is said that ... year after year we have around 980 boys in top 1000 ranks. Generally we see only 4 to 5 girls in top 500. In last 50 years not once any girl topped in IIT-JEE advanced. Forget about Single digit ranks, double digit ranks by girls have been extremely rare. It is all about “ good boys “, “ hard working “, “ focused “, “[Bel-esprit](#) “ **boys**.

In 2015, Only 2.6% of total candidates who qualified are girls (upto around 12,000 rank). while 20% of the Boys, amongst all candidates qualified. The Total number of students who appeared for the exam were around 1.4 million for IIT-JEE main. Subsequently 1.2 lakh (around 120 thousands) appeared for IIT-JEE advanced.

IIT-JEE results and analysis, of many years is given at <https://zookeepersblog.wordpress.com/iit-jee-iseet-main-and-advanced-results/>

In Bangalore it is rare to see a girl with rank better than 1000 in IIT-JEE advanced. We hardly see 6-7 boys with rank better than 1000. Hardly 2-3 boys get a rank better than 500.

See <http://skmclasses.weebly.com/everybody-knows-so-you-should-also-know.html>

Thousands of people are exposing the heinous crimes that Motherly Women are doing, or Female Teachers are committing. See <https://www.facebook.com/WomenCriminals/>

Some Random Examples must be known by all

It is extremely unfortunate that the " woman empowerment " has created. This is the kind of society and women we have now. I and many other sensible Men hate such women. Be away from such women, be aware of reality.



Mother Admits On Facebook to Sleeping with 15 Yr Old Son, They Have a Baby Together - Alwaysturntup

Sometimes it hard to believe w From Alwaysturntup

ALWAYZTURNUP.ME



'Sex with my son is incredible - we're in love and we want a baby'

Ben Ford, who ditched his wife when he met his mother Kim West after 30 years, claims what the couple are doing 'isn't incest'

MIRROR.CO.UK

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

Woman sent to jail for the rest of her life after raping her four grandchildren is described as the 'most evil person' the judge has ever seen

Edwina Louis rape...

[See More](#)



Former Shelbyville ISD teacher who had sex with underage student gets 3 years in prison

After a two day break over the weekend, A Shelby County jury was back in the courtroom looking to conclude the trial of a former Shelbyville ISD teacher who had...

KLTV.COM | BY CALEB BEAMES



Woman sent to jail for raping her four grandchildren

A Ohio grandmother has been sentenced to four consecutive life terms after being found guilty of the rape of her own grandchildren. Edwina Louis, 53, will spend the rest of her life behind bars.

DAILYMAIL.CO.UK

<http://www.thenativecanadian.com/.../eastern-ontario-teacher-...>



The N.C. Chronicles.: Eastern Ontario teacher charged with 36 sexual offences

anti feminism, Child abuse, children's rights, Feminist hypocrisy,

THENATIVECANADIAN.COM | BY BLACKWOLF



Hyd woman kills newborn boy as she wanted daughter - Times of India

Having failed to bear a daughter for the third time, a shopkeeper's wife slit the throat of her 24day-old son with a shaving blade and left him to die in a street on Tuesday night.Purnima's first child was a stillborn boy, followed by another boy born five years ago.

TIMESOFINDIA.INDIATIMES.COM

Montgomery's son, Alan Vonn Webb, took the stand and was a key witness in her conviction.
 "I want to see her placed somewhere she can never do that to children

 See More



Woman sentenced to 40 years in prison for raping her children
 A Murfreesboro mother found guilty of raping her own children learned her fate on Wednesday.
 WAFF.COM | BY DENNIS FERRIER

gentler sex? Violence against men.'s photo.



Women, the gentler sex? Violence against men.
 April 8 at 1:38am · Like Page
 In fact, the past decade has seen a dramatic increase in the number of incidents of women raping and sexually assaulting boys and men. On May 2014, Jezebel repo...

End violence against women



North Carolina Grandma Eats Her Daughter's New Born Baby After Smoking Bath Salts
 Henderson, North Carolina—A North Carolina grandmother of 4 and recovering drug addict, is now in custody after she allegedly ate her daughter's newborn baby....
 AZ-365.TOP



28-Year-Old Texas Teacher Accused of Sending Nude Picture to 14-Year-Old Former Student
 BREITBART.COM

<http://latest.com/.../attractive-girl-gang-lured-men-alleywa.../>



Attractive Girl Gang Lured Men Into Alleyways Where Female Body Builder Would Attack Them
 A Mexican street gang made up entirely of women has been accused of using their feminine wiles to lure men into alleyways and then beating them up and...
 LATEST.COM

http://www.wfmj.com/.../youngstown-woman-convicted-of-raping-...



Youngstown woman convicted of raping a 1 year old is back in jail
 A Youngstown woman who went to prison for raping a 1-year-old boy fifteen years ago is in trouble with the law again.
 WFMJ.COM

End violence against women



Women are raping boys and young men

Rape advocacy has been maligned and twisted into a political agenda controlled by radicalized activists. Tim Patten takes a razor keen and well supported look into the manufactured rape culture and...

AVOICEFORMEN.COM | BY TIM PATTEN



Bronx Woman Convicted of Poisoning and Drowning Her Children

Lisette Bamenga researched methods on the Internet before she killed her son and daughter in 2012.

NYTIMES.COM | BY MARC SANTORA

A Russian-born newlywed slowly butchered her German husband — feeding strips of his flesh to their dog until he took his last breath. Svetlana Batukova, 46, was...

See More



She killed her husband and then fed him to her dog: police

A Russian-born newlywed butchered her German hubby — and fed strips of his flesh to her pooch, authorities said. Svetlana Batukova offered Horst Hans Henkels at their...

NYPOST.COM

April 4 at 4:48am



Female prison officers commit 90pc of sex assaults on male teens in US juvenile detention centres

Lawsuit in Idaho highlights the prevalence of sexual victimization of juvenile offenders.

BTIMES.CO.UK | BY NICOLE ROJAS



Daily Mail

January 15, 2015

Mother charged with rape and sodomy of her son's 12-year-old friend



Mom, 30, 'raped and had oral sex with her son's 12-year-old friend'

Nicole Marie Smith, 30, (pictured) of St Charles County, Missouri, has been jailed after she allegedly targeted the 12-year-old boy at her home.

DAILYM.AI

This mother filmed herself raping her own son and then sold it to a man for \$300. The courts just decide her fate. When you see what she got, you're going to be outraged.



Mother Who Filmed Herself Raping Her 1-Year-Old Son Receives Shocking Sentence

"...then used the money to buy herself a laptop..."

AMERICANNEWS.COM

In several countries or rather in several regions of the world, family system has collapsed, due to bad nature and naughty acts of women. Particularly in Britain, and America, almost 50% people are alone, lonely, separated, divorced or failed marriages. In 2013, 48% children were born out of wedlock. It was projected that by 2016, more than 51% children will be born, to unmarried mothers. In these developed countries " paternity fraud " by women, are close to 20%. You can see several articles in the net, and in wikipedia etc. This means 1 out of 5 children are calling a wrong man as dad. The lonely, alone " mothers " are frustrated. They see the children as burden. Love in the Society in general is lost, long time ago. The types of " Mothers " and " Women " we have now

This is the type of women we have in this world. These kind of women were also someones daughter



Mother Stabs Her Baby 90 Times With Scissors After He Bit Her While Breastfeeding Him!

Eight-month-old Xiao Bao was discovered by his uncle in a pool of blood. He needed 100 stitches after the incident, he is now recovering in hospital. Reports say his...

MOMMABUZZ.COM





HURT FEMINISM BY DOING NOTHING

- ✗ DON'T HELP WOMEN
- ✗ DON'T FIX THINGS FOR WOMEN
- ✗ DON'T SUPPORT WOMEN'S ISSUES
- ✗ DON'T COME TO WOMEN'S DEFENSE¹
- ✗ DON'T SPEAK FOR WOMEN
- ✗ DON'T VALUE WOMEN'S FEELINGS
- ✗ DON'T PORTRAY WOMEN AS VICTIMS
- ✗ DON'T PROTECT WOMEN²

✓ WITHOUT WHITE KNIGHTS
 FEMINISM WOULD END TODAY

¹Don't even nawalt ("Not All Women Are Like That") ²for example from criticism or insults


How Society prioritize Men



Professor Subhashish Chattopadhyay

Spoon Feeding Series - Reduction Methods in Organic Chemistry

We must first understand the most Important graph or Image of Chemistry



It is very strange, that so many Chemistry books exist in this world, which do not even have a single Periodic table in them. **Almost all books do not have this graph either.** This is the graph of electrode potential of various elements with respect to (standard) Hydrogen electrode.

All chemical reactions happen because of this electrode potential difference. When a Solid is added to a solution of another compound such as a Gold Spoon in CuSO_4 Solution, or a Copper container having AgNO_3 Solution, then the displacement reaction happens or not is decided by these potential values. Battery and Electrochemistry chapter uses this knowledge. Oxidation-Reduction reactions, titrations etc all need this understanding.

The e.m.f of the standard cell $\text{Zn} | \text{Zn}^{2+} || \text{Ag}^+ | \text{Ag}$ $E^\circ_{\text{cell}} = 1.562$ Volt

If the standard electrode potential of Zinc is -0.762 Volt, then that of Silver is ?

Solution : $E^\circ_{\text{cell}} = E_{\text{Right}} - E_{\text{Left}} = E_{\text{Ag}^+ | \text{Ag}} - E_{\text{Zn}^{2+} | \text{Zn}} = (E_{\text{Reduction (Right)}} - E_{\text{Reduction (Left)}})$

$$\text{So } 1.562 = E(\text{Ag}^+ | \text{Ag}) - (-0.762) \Rightarrow E_{\text{Ag}^+ | \text{Ag}} = 0.8 \text{ Volt}$$

What will happen if a solution of copper sulphate put into an iron pot ?

Note : Electrode potential of Cu is more positive (so higher) than Fe (Iron).

A solution of copper sulphate put into an iron pot will damage. This is because the standard electrode potential for Fe^{2+}/Fe is more negative than that for Cu^{2+}/Cu . The iron oxidises first, while copper ions are reduced. Iron will corrode (rust). Or in more simple words it is easier to replace Cu ions by Fe ions from the pot. Any element lower in the series tends to replace the element which is higher up the series.

This is called a redox reaction, since the iron is being oxidised, and copper ions reduced.

Note : A solution of Iron Salt in a Copper Vessel would have just done fine. In this case copper is in solid form, as vessel. So will not replace the Iron ions.

Half-Reaction	E (V)
$\text{F}_2(\text{g}) + 2\text{e}^- \rightarrow 2\text{F}^-(\text{aq})$	+2.87
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{MnO}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{Mn}^{2+}(\text{aq}) + 2\text{H}_2\text{O}(\text{l})$	+1.23
$\text{NO}_3^-(\text{aq}) + 4\text{H}^+(\text{aq}) + 3\text{e}^- \rightarrow \text{NO}(\text{g}) + 2\text{H}_2\text{O}(\text{l})$	+0.96
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{g}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightarrow 4\text{OH}^-(\text{aq})$	+0.40
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0.00
$\text{N}_2(\text{g}) + 5\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow \text{N}_2\text{H}_5^+(\text{aq})$	-0.23
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.05

↑ strength of oxidizing agent

↓ strength of reducing agent

A more detailed table


Standard Reduction Potentials at 25°C (298 K) for Many Common Half-Reactions			
Half-Reaction	E° (V)	Half-Reaction	E° (V)
$F_2 + 2e^- \rightarrow 2F^-$	2.87	$O_2 + 2H_2O + 4e^- \rightarrow 4OH^-$	0.40
$Ag^{2+} + e^- \rightarrow Ag^+$	1.99	$Cu^{2+} + 2e^- \rightarrow Cu$	0.34
$Co^{3+} + e^- \rightarrow Co^{2+}$	1.82	$Hg_2Cl_2 + 2e^- \rightarrow 2Hg + 2Cl^-$	0.27
$H_2O_2 + 2H^+ + 2e^- \rightarrow 2H_2O$	1.78	$AgCl + e^- \rightarrow Ag + Cl^-$	0.22
$Ce^{4+} + e^- \rightarrow Ce^{3+}$	1.70	$SO_4^{2-} + 4H^+ + 2e^- \rightarrow H_2SO_3 + H_2O$	0.20
$PbO_2 + 4H^+ + SO_4^{2-} + 2e^- \rightarrow PbSO_4 + 2H_2O$	1.69	$Cu^{2+} + e^- \rightarrow Cu^+$	0.16
$MnO_4^- + 4H^+ + 3e^- \rightarrow MnO_2 + 2H_2O$	1.68	$2H^+ + 2e^- \rightarrow H_2$	0.00
$2e^- + 2H^+ + IO_4^- \rightarrow IO_3^- + H_2O$	1.60	$Fe^{3+} + 3e^- \rightarrow Fe$	-0.036
$MnO_4^- + 8H^+ + 5e^- \rightarrow Mn^{2+} + 4H_2O$	1.51	$Pb^{2+} + 2e^- \rightarrow Pb$	-0.13
$Au^{3+} + 3e^- \rightarrow Au$	1.50	$Sn^{2+} + 2e^- \rightarrow Sn$	-0.14
$PbO_2 + 4H^+ + 2e^- \rightarrow Pb^{2+} + 2H_2O$	1.46	$Ni^{2+} + 2e^- \rightarrow Ni$	-0.23
$Cl_2 + 2e^- \rightarrow 2Cl^-$	1.36	$PbSO_4 + 2e^- \rightarrow Pb + SO_4^{2-}$	-0.35
$Cr_2O_7^{2-} + 14H^+ + 6e^- \rightarrow 2Cr^{3+} + 7H_2O$	1.33	$Cd^{2+} + 2e^- \rightarrow Cd$	-0.40
$O_2 + 4H^+ + 4e^- \rightarrow 2H_2O$	1.23	$Fe^{2+} + 2e^- \rightarrow Fe$	-0.44
$MnO_2 + 4H^+ + 2e^- \rightarrow Mn^{2+} + 2H_2O$	1.21	$Cr^{3+} + e^- \rightarrow Cr^{2+}$	-0.50
$IO_3^- + 6H^+ + 5e^- \rightarrow \frac{1}{2}I_2 + 3H_2O$	1.20	$Cr^{3+} + 3e^- \rightarrow Cr$	-0.73
$Br_2 + 2e^- \rightarrow 2Br^-$	1.09	$Zn^{2+} + 2e^- \rightarrow Zn$	-0.76
$VO_2^+ + 2H^+ + e^- \rightarrow VO^{2+} + H_2O$	1.00	$2H_2O + 2e^- \rightarrow H_2 + 2OH^-$	-0.83
$AuCl_4^- + 3e^- \rightarrow Au + 4Cl^-$	0.99	$Mn^{2+} + 2e^- \rightarrow Mn$	-1.18
$NO_3^- + 4H^+ + 3e^- \rightarrow NO + 2H_2O$	0.96	$Al^{3+} + 3e^- \rightarrow Al$	-1.66
$ClO_2 + e^- \rightarrow ClO_2^-$	0.954	$H_2 + 2e^- \rightarrow 2H^-$	-2.23
$2Hg^{2+} + 2e^- \rightarrow Hg_2^{2+}$	0.91	$Mg^{2+} + 2e^- \rightarrow Mg$	-2.37
$Ag^+ + e^- \rightarrow Ag$	0.80	$La^{3+} + 3e^- \rightarrow La$	-2.37
$Hg_2^{2+} + 2e^- \rightarrow 2Hg$	0.80	$Na^+ + e^- \rightarrow Na$	-2.71
$Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.77	$Ca^{2+} + 2e^- \rightarrow Ca$	-2.76
$O_2 + 2H^+ + 2e^- \rightarrow H_2O_2$	0.68	$Ba^{2+} + 2e^- \rightarrow Ba$	-2.90
$MnO_4^- + e^- \rightarrow MnO_4^{2-}$	0.56	$K^+ + e^- \rightarrow K$	-2.92
$I_2 + 2e^- \rightarrow 2I^-$	0.54	$Li^+ + e^- \rightarrow Li$	-3.05
$Cu^+ + e^- \rightarrow Cu$	0.52		


Spoon Feeding

What will happen if a solution of Magnesium sulphate put into an Copper pot ?

Note : Magnesium loses electron very easily, metallic character, or basic character being high. Electrode potential is - 2.37 While we know Copper has +ve electrode potential. (This is an important exception and the data is better remembered as 0.5)

So Copper is solid in Copper pot (or even if a copper spoon was inserted, the same logic). Copper will not replace Magnesium ions. But if a Magnesium spoon was put into Copper-Sulphate solution then the Magnesium ions / atoms will start replacing the Cu ions and Copper will start getting precipitated. The Magnesium spoon, or container will start getting corroded or damaged.






Nernst Equation also uses Electrode potential

Walther Hermann Nernst (1864 - 1941) German Physicist & Chemist received the Nobel prize in 1920 Chemistry.



$$\text{At } 25^{\circ}\text{C}; E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.059}{n} \log \frac{M}{[M^{n+}]} \text{ or } \frac{1}{[M^{n+}]}$$

$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.059}{n} \log \left[\frac{\text{Products}}{\text{Reactants}} \right] \text{ or } Q.$$

$$E_{\text{cell}} = E_{\text{cathode}}^0 - E_{\text{anode}}^0 - \frac{0.059}{n} \log \left[\frac{P}{R} \right]$$



Consider the following set of half-reactions



This is valid only if all species have concentrations of 1 M.

Le Châtelier's principle gives us some ideas. Look at the overall reaction. Consider the reaction to be at equilibrium ($E_{\text{cell}} = 0 \text{ V}$). If we add more Fe^{2+} to the reaction mixture, The equilibrium will shift to the right to use up the excess Fe^{2+} . This will, of course cause a positive cell potential to be measurable.

Conversely, if we removed Fe^{2+} from the reaction mixture, the equilibrium would shift to the left to try to replace some of the missing Fe^{2+} . This would result in a measurable negative potential (reaction goes left).

If the reaction were not at equilibrium then these preceding two changes would be additive to the measured voltage (say standard voltage). So, if we add Fe^{2+} to a mixture at Standard Conditions, then the voltage would be more positive and if we removed some Fe^{2+} from a mixture at Standard Conditions the measured voltage would be lower (less positive).

The reverse results would be found for addition or removal of Zn^{2+} (a product). More Zn^{2+} would give a lower positive potential and *vice versa*.

Recall the relation: $\Delta G = \Delta G^\circ + RT \ln Q$ (for non-standard conditions)

Substitute the new definition of ΔG and we get

$$-nFE = -nFE^\circ + RT \ln Q$$

$$E = E^\circ - \frac{RT}{nF} \ln Q$$

(This is called the *Nernst Equation*).

Thus, we can calculate the cell potential of any cell, given only the initial conditions (to calculate Q) and the reduction potentials (to calculate E°)

We can calculate the voltage for the situation where the concentrations are not standard. For example, consider $[\text{Fe}^{2+}] = 0.1 \text{ M}$ and $[\text{Zn}^{2+}] = 1.9 \text{ M}$

For this reaction, we also know that there are two electrons transferred for each equation ($n = 2$).






$$E = E^\circ - \frac{RT}{nF} \times \ln \frac{[Zn^{2+}]}{[Fe^{2+}]}$$

$$= 0.32V - \frac{8.3145 \frac{J}{mol \cdot K} \times 298.15K}{2 \times 96485 \frac{C}{mol}} \times \ln \frac{1.9}{0.10}$$

$$= 0.28V$$

Although the reaction is 90% complete (from Standard Conditions initial) the cell potential has only dropped by a small amount (0.04 V). This is normal. For example, the batteries in your flashlight will put out almost full voltage until the last traces of chemical are almost used up at which point the voltage will drop off rather sharply. This is an especially useful characteristic of cells powering electronic equipment.


A voltaic cell using inactive electrodes.



↑
The inert electrode is specified.


↑
A comma is used to show components that are in the same phase.

A voltaic cell is made of aqueous Bromine and Zinc. Given -



Calculate E_{bromine} , given that $E_{\text{zinc}} = -0.76 \text{ V}$

Solution :





$$E_{\text{cell}} = E_{\text{cathode}} - E_{\text{anode}}$$

$$1.83 = E_{\text{bromine}} - (-0.76)$$

$$1.83 - 0.76 = E_{\text{bromine}} \quad = 1.07 \text{ Volt}$$


What is the equilibrium constant for the reaction of copper metal with bromine to form copper(II) ions and bromide ions in aqueous solution at 25°C?



$$\ln K = \frac{nFE^\circ}{RT} = \frac{2 \times 96458 \frac{\text{C}}{\text{mol}} \times 0.75 \text{ V}}{8.31451 \frac{\text{J}}{\text{mol K}} \times 298.15 \text{ K}} = 58.38_{208}$$


$$K = e^{58.38_{208}} = 2.2 \times 10^{25}$$

Using half reactions write the spontaneous Redox reactions




We have to multiply the Silver reaction with 2 to balance and match electrons

Note : This does not change the voltage obtained



Now we need to add the half reactions to eliminate the electrons, but subtract to get the Voltage correctly. Recall $E^\circ_{\text{cell}} = E_{\text{Right}} - E_{\text{Left}} = E_{\text{Cathode}} - E_{\text{Anode}} = (E_{\text{Reduction (Right)}} - E_{\text{Reduction (Left)}})$




$$E^\circ_{\text{cell}} = E^\circ_{\text{silver}} - E^\circ_{\text{tin}} = 0.80 - (-0.14) = 0.94 \text{ V}$$

Cathode provides electrons. Ag ions are receiving electrons at Cathode. Anode is positive where electrons are lost. See Tin is more metallic (than Silver) and is loses electrons in Anode.

It is easier to visualize the Voltage as gap in electrode potential. Silver is higher up than Hydrogen in the series. Thus has positive electrode potential. While Tin is more metallic, below Hydrogen in the series. The gap is of 0.94 Volts.


-

Zinc and Magnesium Oxide Battery



-


Zinc and Magnesium Oxide Battery




The mercury battery uses HgO as the oxidizing agent instead of Ag₂O and has cell potential of 1.3 V.

-


The Lithium Batteries use Silver Vanadium Pentoxide as cathode




The most commonly used Battery is Lead-Acid Battery



Cathode (reduction):



Overall (cell) reaction (discharge):





Overall (cell) reaction (recharge):




-

Nickel Metal Hydride Battery




Overall (cell) reaction:




-

Laptops, Cell-phones most commonly use Lithium ion Batteries


Anode (oxidation):



Cathode (reduction):





Overall (cell) reaction:



-


Hydrogen Fuel Cell



-

What is the value of the solubility product constant for AgCl?

We could calculate this by breaking the overall K_{sp} reaction into a series of redox reactions as follows:



$$\ln K = \frac{nFE^\circ}{RT} = \frac{1 \times 96485 \frac{\text{C}}{\text{mol}} (-0.58 \text{ V})}{8.3145 \frac{\text{J}}{\text{mol K}} \times 298.15 \text{ K}} = 22.5744$$

$$K = 6.3 \times 10^{-9}$$

-

Ion Concentrations can also be calculated...

Consider the Platinum-Hydrogen electrode coupled with a copper/copper(II) electrode.

The overall reaction is $\text{Cu}^{2+} + \text{H}_2 \rightarrow \text{Cu} + 2\text{H}^+$.

The Nernst equation is

$$E = E^\circ - \frac{8.31451 \frac{\text{J}}{\text{mol K}} \times 298.15 \text{ K}}{2 \text{ mol} \times 96485 \frac{\text{C}}{\text{mol}}} \times \ln \left(\frac{a(\text{H}^+)^2}{a(\text{Cu}^{2+}) \times a(\text{H}_2)} \right)$$

$$E = E^\circ - \frac{.0257 \text{ V}}{2} \times \ln \left(\frac{a(\text{H}^+)^2}{a(\text{Cu}^{2+}) \times a(\text{H}_2)} \right) \quad \text{since}$$

$$\frac{8.31451 \frac{\text{J}}{\text{mol K}} \times 298.15 \text{ K}}{96485 \frac{\text{C}}{\text{mol}}} = .0256928 \text{ V}$$

If $[\text{Cu}^{2+}] = 1 \text{ M}$ and $P(\text{H}_2) = 1 \text{ bar}$ (both their activities are 1) then

$$E = E^\circ - \frac{0.0257 \text{ V}}{2} \ln [\text{H}^+]^2$$

$$E = E^\circ - 0.0257 \text{ V} \ln [\text{H}^+] \quad (\text{Note: } \ln Y = 2.303 \times \log Y \text{ and } -.257 \times 2.303 = .05917)$$


$$E = E^\circ + 0.05917 \text{ V} (-\log [\text{H}^+])$$

$$E = E^\circ + 0.05917 \text{ V} \times \text{pH}.$$

We see here that the cell potential is a function of pH. The probes in pH meters are set up this way. A complete electrochemical cell is contained within the probe casing. All chemicals are at standard conditions and a porous glass membrane allows only H^+ ions to pass through.

Electrochemical Cell

The electrochemical cell shown below is a concentration cell.




The emf of the cell depends on the difference in concentration of M^{2+} ions at the two electrodes. The emf of the cell at 298 K is 0.059 V.

- Q The solubility product (K_{sp}) of MX_2 at 298 K based on the information available for the given concentration cell (take $2.303 RT/F = 0.059 \text{ V}$ at 298 K) is
- (a) $1 \times 10^{-15} \text{ mol}^3 \text{ dm}^{-9}$ (b) $4 \times 10^{-15} \text{ mol}^3 \text{ dm}^{-9}$
(c) $1 \times 10^{-12} \text{ mol}^3 \text{ dm}^{-9}$ (d) $4 \times 10^{-12} \text{ mol}^3 \text{ dm}^{-9}$
- Q The value of ΔG for the given cell reaction (take $1F = 96500 \text{ C mol}^{-1}$) is
- (a) -5.7 kJ mol^{-1} (b) 5.7 kJ mol^{-1} (c) 11.4 kJ mol^{-1} (d) $-11.4 \text{ kJ mol}^{-1}$

Solutions :


The cell reaction may be obtained as follows.



The cell potential is

$$E = - \frac{RT}{2F} \ln \left\{ \frac{[\text{M}^{2+}]_{\text{L}} / \text{mol dm}^{-3}}{[\text{M}^{2+}]_{\text{R}} / \text{mol dm}^{-3}} \right\} \quad \text{i.e.,} \quad 0.059 \text{ V} = - \left(\frac{0.059 \text{ V}}{2} \right) \log \left\{ \frac{[\text{M}^{2+}]_{\text{L}} / \text{mol dm}^{-3}}{0.001} \right\}$$

This gives $\log ([\text{M}^{2+}]_{\text{L}} / \text{mol dm}^{-3}) = -2 + \log (0.001) = -2 - 3 = -5$ i.e., $[\text{M}^{2+}]_{\text{L}} = 1.0 \times 10^{-5} \text{ mol dm}^{-3}$
From the reaction



The ΔG of the cell reaction is

$$\Delta G = - nFE = - (2) (96500 \text{ C mol}^{-1}) (0.059 \text{ V}) = - 11387 \text{ J mol}^{-1} = - 11.4 \text{ kJ mol}^{-1}$$

Hence, we have

The **choice (b)** is correct in first Question

The **choice (d)** is correct in the second Question

Reduction Methods in Organic Chemistry

Let us always remember the basics. If something, say M, loses electrons, it is getting oxidized. Metals are basic. Metals lose electrons and get oxidized. The metals which lose electron very easily, get oxidized very easily. So these are better reducing agents.

Adding of Oxygen, or Halogens (F, Cl, Br, I) or other p-Block elements which accept electrons, is oxidation. So if Potassium (K) gets Oxygen (attaches to O) to form K_2O or KO_2 (Superoxide), K is getting oxidized. If Cs attaches with F then Cs is getting oxidized, while F is being reduced. Cs is a reducing agent as it is losing electrons. F is an oxidizing agent as it is gaining electrons.


Adding of Hydrogen is reduction. So if Ethene receives Hydrogen to become Ethane, then Ethene is being reduced. Raney Nickel is Nickel with molecular level pores. Raney Ni with HCl gives Hydrogen atoms, (some teachers say this as Nascent Hydrogen) locally, meaning near the molecule where the reaction is needed. So Raney Ni + HCl is a good reducing agent. Same for Zinc dust + HCl. But then what is Birch reduction ?

As my experience of teaching more than 25 years, almost every student understands these. Knowing this is essential but does not ensure success in Organic chemistry. There are many kinds of details of oxidizing agent or Reducing agent that vary from reactions / situations.

I reasonably searched and realized that details of Oxidation or Reduction is scattered. There is no single e-Book that gives all of these as a collection.


So this e-Book is free for you. All information at a single place.

Gilman's Reagent (An organometallic compound)




Br or Oxygen atoms were being knocked out. Removal of Oxygen or halogen is reduction.

When an aldehyde is to be prepared from an alkyl cyanide, the cyanide has to be partially reduced with SnCl_2/HCl (Stephen's reduction) and when ketone is to be formed, the cyanide is treated with Grignard reagent.




Adding nascent hydrogen is Reduction.



RMgI is Grignard reagent. R always becomes Anion, i.e. negatively charged ion. MgI is cation, i.e. positively charged. With water OH group joins Mg.


The reducing action of sodium borohydride is so specific that it reduces only $\text{C}=\text{O}$ bond but not $\text{C}=\text{C}$ bond.

For example :



NaBH_4 is a strong reducing agent. Aldehyde in this case reduces to alcohol.

Preparation of ethanal from carbon and hydrogen can be effected by following reaction :



The carbonyl group $\left(\text{>C}=\text{O} \right)$ can be reduced to methy-


lene group $\left(\text{>CH}_2 \right)$ resulting in the formation of alkanes by

any of the following reagents :

- (i) Zn-Hg/HCl : Clemmensen's reduction
- (ii) $\text{H}_2\text{N-NH}_2/\text{KOH}$: Wolff kishner's reduction
- (iii) HI/P at 424 K

This was an industrial way of adding hydrogen. (thus reduction).


Trimethyl silyl iodide:




Peterson's synthesis: Beta hydroxyl Trimethyl silane

Silyl wittig reaction

Carbonyl compounds \longrightarrow Alifins



H₂/Ni :



Baker's yeast:



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

RMgX :

Acid derivatives + 1 eq RMgX \longrightarrow Carbonyl compounds

Acid derivatives + 2 eq RMgX \longrightarrow 3^o Alcoholic compounds

Tri n-Butyl tin hydride(nBu₃SnH):

In presence of thermal/light/peroxides

R-F < R-Cl < R-Br < R-I

R-X \longrightarrow R-H




Wolf Kishner reduction (Knocking out Oxygen)

Wolf kishner reduction(H₂N-NH₂/KOH)



LDA(Lithium di isopropyl amide):


Strong base due to 2^o amine & free e⁻ pair. Sterically hindered base



Wilkinson's catalyst[Tris (tri phenyl phosphine)rhodium(I)chloride]:

Hydrogenation of C=C, C≡C, with out disturbing other functional groups,

It adds H₂ to olefins at same side, Cis olefin to meso, Trans olefin to racemic mixture..




Pi bonds were being broken and hydrogen atoms were being added.

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

Palladium/H₂ :




H₂/Ni useful application:




Birch Reduction is the only way to break the double bonds of Benzene and convert to cyclohexane. Benzene ring is pretty stable, and known as resonance stabilization. Adding hydrogen (i.e. reduction) of benzene is not easy.

Birch reduction:




Metal surface catalysts (Palladium in this case) gives cis isomer. This is because an intermediate is formed where both sides join the metal surface. Both sides being towards the metal catalyst eventually makes it cis.

Pd-BaSO₄(Lindlar's catalyst):




Benkeser reduction:



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

Clemmenson's (Zn-Hg/HCl):




Clemmenson's reduction is Zn / Hg with HCl


Alanes(AlH_3):



With silanes:




Reduction with HCO_2H :




Sodium boro hydride:


Reduces predominately unconjugated carbonyl compounds




Vils meier reaction:



Silver tetra fluoro borate(AgBF_4):



Luche's reagent ($\text{NaBH}_4 \cdot \text{CeCl}_3$): Reducing conjugating carbonyl compounds,
Reduces ketones more efficiently than aldehydes




Super hydride (LiBHET_3):



Sodium cyano boro hydride (NaBH_3CN):


Readily reduces iodides, bromides, and tosylates to the hydrocarbons...

HMPA= hexa methyl phosphoramide




X=I, Br, OTs


Ketones reacts with Grignard reagents giving tertiary alcohols.



or




Example:



DIBAL-H (Di iso butyl aluminium hydride):

High temperatures: Acid derivatives to Alcohols


Low temperatures: Acid derivatives to Aldehydes



Adams catalyst ($\text{H}_2\text{-PtO}_2$): Less hindered alkenes to alkanes.




Rosenmund reduction:




Grignard reactions of carbonyl compounds


Formaldehyde ($\text{H}_2\text{C=O}$) reacts with Grignard reagents giving primary alcohol.




or




Example:




Aldehydes reacts with Grignard reagents giving secondary alcohols.




or




Example:



- On adding water, hydrolysis of the intermediate yields the primary alcohols.




- LiAlH_4 has no effect on the benzene ring or the double bond.
- $-\text{COOH}$ is reduced to $-\text{CH}_2\text{OH}$ but the $\text{C}=\text{C}$ bonds remains unchanged.




NaNH_2 is a strong base and excellent nucleophile. It's used for deprotonation of weak acids and also for elimination reactions. Similar to: LDA (lithium diisopropylamide). The bases LiNH_2 and KNH_2 essentially similar. As a strong base, NaNH_2 will deprotonate alkynes, alcohols, and a host of other functional groups with acidic protons such as esters and ketones. It is also a very strong nucleophile. One common application of NaNH_2 is in the deprotonation of alkynes to give so-called "acetylide" ions. These ions are excellent nucleophiles and can go on to react with carbonyls in addition reactions.

Deprotonation of alkynes




A second application of NaNH_2 is in the formation of alkynes from halogens. Treatment of either geminal dihalides (i.e. - two halogens on one carbon) or vicinal dihalides (halogens on adjacent carbons) with NaNH_2 (2 equiv) will result in the formation of alkynes. So this is reducing property.

Formation of alkynes from geminal dihalides




Formation of alkynes from vicinal dihalides




Since vicinal dihalides are easily made by the reaction of alkenes with halogens such as Br₂ or I₂, this is a useful way of converting alkenes to alkynes.


Deprotonation of functional groups such as OH and even alkyne C-H should hopefully be straightforward, but the use of bases to make alkenes may require some explanation. This is what is known as an elimination reaction, in that the elements H and Br (in this example) are removed in order to form the alkene. Specifically, this is an example of an E2 reaction.




Since the alkene still has a halide attached, this too can be removed to generate a second double bond (π bond). This is another example of the E2 in that the hydrogen has to be anti to the bromine that is eliminated, but is unusual in that it is an sp² hydrogen that is affected here:





NaNH₂ can replace Chlorine




Furthermore, since NaNH₂ is a strong base, it has the significant disadvantage of promoting side reactions from elimination (this can occur when attempting an S_N2 with NaNH₂ as the nucleophile, for example). Therefore, it is generally wise to avoid using NaNH₂ as a nucleophile in organic synthesis. Sodium azide (followed by reduction) is the usual substitute.



NaNH₂ acts as reducing agent




Benzyne Reaction: Elimination- Addition



Reactant is halobenzene with no electron-withdrawing groups on the ring.

Use a very strong base like NaNH_2 .

Another reaction done in cold



Topic wise grouping of information kind of Dictionary of General Chemistry

Graphite and Diamond comparison

With respect to graphite and diamond, which of the following statement(s) given below is(are) correct?

- (a) Graphite is harder than diamond.
- (b) Graphite has higher electrical conductivity than diamond.
- (c) Graphite has higher thermal conductivity than diamond.
- (d) Graphite has higher C—C bond order than diamond.

Solution :

The given facts about graphite and diamond are as follows.

Graphite is less hard than diamond. Graphite has higher electrical and thermal conductivities than diamond. This is due to highly delocalized nature of π electrons. Graphite has higher C—C bond order (due to σ and π bonds) than that of diamond (which has only σ bonds)

Therefore, the **choices (b), (c) and (d)** are correct.

Stability of lyophobic colloidal particles

Choose the correct reason(s) for the stability of lyophobic colloidal particles.

- (a) Preferential adsorption of ions on their surface from the solution.
- (b) Preferential adsorption of solvent on their surface from the solution.
- (c) Attraction between different particles having opposite charges on their surface.
- (d) Potential difference between the fixed layer and the diffused layer of opposite charges around the colloidal particles.

Solution :



The stability of lyophobic colloidal particles is due to the preferential adsorption of common ion from the solution. Each colloidal particle has the same charge and are thus kept away from each other due to electronic repulsion (choice a).



There exists potential difference between the fixed and diffused layer of opposite charge (choice d).

Therefore, the **choice (a) and (d)** are correct.

Physiorption and Chemisorption comparison

The given graph/data I, II, III and IV represent general trends observed for different physisorption and chemisorption processes under mild conditions of temperature and pressure. Which of the following choice(s) about I, II, III and IV is(are) correct.





- (a) I is physisorption and II is chemisorption
(b) I is physisorption and III is chemisorption
(c) IV is chemisorption and II is chemisorption
(d) IV is chemisorption and III is chemisorption

Solution:

We have the following facts.

- Adsorption decreases with increase in temperature at constant pressure in case of physisorption
- Adsorption increases initially with increase in temperature at constant pressure in case of chemisorption as more molecules satisfy the energy of activation.
- The enthalpy change during chemisorptions is high due to the bond formation between adsorbent and adsorbate.

With these guidelines, we will have

Graph I represents physisorption


Graph III represents physisorption

Graph II represents chemisorption

Graph IV represents chemisorptions

Therefore, The **choices (a) and (c)** are correct.

A compound M_pX_q has cubic close packing (CCP) arrangement of X. Its unit cell structure is shown below.



The empirical formula of the compound is

- (a) MX (b) MX_2 (c) M_2X (d) M_5X_{14}

Solution :

The atom at the corner of the cube is shared amongst 8 unit cells; the atom at the centre of a face is shared between 2 unit cells, the atom at the centre of edges is shared amongst 4 unit cells and that at the centre of the cube belongs wholly to the unit cell.

There are eight X atoms at the corners and six X atoms at the centre of faces. Thus, the number of X atoms per

unit cell is

$$8\left(\frac{1}{8}\right) + 6\left(\frac{1}{2}\right) = 1 + 3 = 4$$

There are four M atoms at the centre of edges and one M atom at the centre of the cube.
Thus, the number of M atoms per unit cell is

$$4\left(\frac{1}{4}\right) + 1(1) = 1 + 1 = 2$$

The formula of the compound will be M_2X_4 and the empirical formula will be MX_2 .
Therefore, the **choice (b)** is correct.



HCl HBr and HI react with $AgNO_3$ to give a precipitate that dissolves in $Na_2S_2O_3$

Which of the following halides react(s) with $AgNO_3(aq)$ to give a precipitate that dissolves in $Na_2S_2O_3(aq)$?

- (a) HCl (b) HF (c) HBr (d) HI

Solution :

The precipitates of $AgCl$, $AgBr$ and AgI are soluble in $Na_2S_2O_3(aq)$ forming the complex $Na_3[Ag(S_2O_3)_2]$. For example,



AgF is soluble in water.


Therefore, the **choices (a), (c) and (d)** are correct.

Reversible expansion of an ideal gas

The reversible expansion of an ideal gas under adiabatic and isothermal conditions is shown in the figure.

Which of the following statement(s) is(are) correct?

- (a) $T_1 = T_2$
(b) $T_2 > T_1$
(c) $w_{\text{isothermal}} > w_{\text{adiabatic}}$
(d) $\Delta U_{\text{isothermal}} > \Delta U_{\text{adiabatic}}$



Solution :

We will have

$T_1 = T_2$ as the expansion is isothermal.

$T_3 < T_1$ as the adiabatic expansion involves cooling.

$|w_{\text{isothermal}}| > |w_{\text{adiabatic}}|$ as the area under the isothermal curve is greater than that under adiabatic curve.

$\Delta U_{\text{isothermal}} = 0$ as temperature remains constant while $\Delta U_{\text{adiabatic}} < 0$ as the expansion occurs at the expense of internal energy.

Therefore, the **choices (a) and (d)** are correct.

Note: The choice (c) is correct if the magnitude of the work is considered. Since expansion carries negative sign as per IUPAC recommendations, the choice (c) will not be correct if the negative sign is also considered.



Acid Ionization Constant	$K_a = \frac{[H^+][A^-]}{[HA]}$
Adiabatic change	$PV = k$
Base Ionization Constant	$K_b = \frac{[OH^-][HB^+]}{[B]}$
Bohr Radius	$a_0 = \frac{\hbar^2}{m_e k e^2}$
Boiling Point Elevation	$\Delta T_b = i K_b \times \text{molality}$
Buffer Design Equation	$pH \approx pK_a - \log \frac{[HA]_0}{[A^-]_0}$
Cell Voltage	$E_{cell} = E_{cell}^\circ - \frac{RT}{n\mathfrak{F}} \ln Q = E_{cell}^\circ - \frac{0.0592}{n} \log Q$
Charles' Law	$\frac{V}{t} = k$

Atomic Structure

DESCRIPTION	EQUATION
Bohr Radius	$a_0 = \frac{\hbar^2}{m_e k e^2}$
De Broglie Wavelength	$\lambda = \frac{h}{mv}$
Linear Momentum	$p = mv$
Planck's Quantized (Quantum) Energy Equation	$\Delta E = h\nu$
Radii of stable orbits in the Bohr model	$r = n^2 \frac{\hbar^2}{m_e k Z e^2} = n^2 \frac{a_0}{Z}$
Relationship between Energy and Principal Quantum Number	$E_n = -R_H \left(\frac{1}{n^2} \right) = \frac{-2.178 \times 10^{-18}}{n^2} \text{joule}$

Rydberg Equation

$$\Delta E = R_H \left(\frac{1}{n_i^2} - \frac{1}{n_f^2} \right)$$

Speed of Light to Wavelength and
Frequency Relationship

$$c = \lambda \nu$$

Electrochemistry

DESCRIPTION	EQUATION
Cell Voltage	$E_{cell} = E_{cell}^{\circ} - \frac{RT}{n\mathcal{F}} \ln Q = E_{cell}^{\circ} - \frac{0.0592}{n} \log Q$
Electric Current	$I = \frac{q}{t}$
Reaction Quotient	$Q = \frac{[C]^c [D]^d}{[A]^a [B]^b}$ where $aA + bB \rightarrow cC + dD$
Relationship between Equilibrium Constant and Cell Voltage	$\log K = \frac{nE^{\circ}}{0.0592}$

Equilibrium

DESCRIPTION	EQUATION
Acid Ionization Constant	$K_a = \frac{[H^+][A^-]}{[HA]}$
Base Ionization Constant	$K_b = \frac{[OH^-][HB^+]}{[B]}$
Buffer Design Equation	$pH \approx pK_a - \log \frac{[HA]_0}{[A^-]_0}$
Gas Pressure and Concentration Relationship	$K_p = K_c (RT)^{\Delta n}$
Ion Product Constant for Water	$K_w = [OH^-][H^+] = K_a \times K_b$ $= 1.0 \times 10^{-14}$ at $25^{\circ}C$
pH and pOH Relationship	$14 = pH + pOH$
pH Defined	$pH = -\log [H^+]$
pK_a Definition	$pK_a = -\log K_a$

pK_b Definition

$$pK_b = -\log K_b$$

pOH and Base Ionization Equilibrium Constant Relationship

$$pOH = pK_b + \log \frac{[HB^+]}{[B]}$$

Gases, Liquids, and Solutions

DESCRIPTION	EQUATION
Adiabatic change	$PV = k$
Boiling Point Elevation	$\Delta T_b = iK_b \times \text{molality}$
Charles' Law	$\frac{V}{t} = k$
Combined Gas Law	$\frac{P_1V_1}{n_1T_1} = \frac{P_2V_2}{n_2T_2}$
Density of a Material	$D = \frac{m}{V}$
Freezing Point Depression	$\Delta T_f = iK_f \times \text{molality}$
Graham's Law of Effusion	$\frac{r_1}{r_2} = \sqrt{\frac{M_2}{M_1}}$
Ideal gas equation	$PV = nRT$
Kinetic Energy per Mole	$\frac{KE}{\text{mole}} = \frac{3}{2}RTn$
Kinetic Energy per Mole	$\frac{KE}{\text{mole}} = \frac{3}{2}RTn$

Thermochemistry

DESCRIPTION	EQUATION
Acid Ionization Constant	$K_a = \frac{[H^+][A^-]}{[HA]}$
Adiabatic change	$PV = k$
Base Ionization Constant	$K_b = \frac{[OH^-][HB^+]}{[B]}$

Bohr Radius
$$a_0 = \frac{\hbar^2}{m_e k e^2}$$

Boiling Point Elevation
$$\Delta T_b = i K_b \times \text{molality}$$


Buffer Design Equation
$$pH \approx pK_a - \log \frac{[HA]_0}{[A^-]_0}$$

Cell Voltage
$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{n\mathcal{F}} \ln Q = E_{\text{cell}}^\circ - \frac{0.0592}{n} \log Q$$

Charles' Law
$$\frac{V}{t} = k$$

Vander waal's force

For one mole of a van der Waals gas when $b = 0$ and $T = 300$ K, the plot of pV versus $1/V$ is shown below.



The value of the van der Waals constant a is

- (a) $1.0 \text{ L}^2 \text{ atm mol}^{-2}$ (b) $4.5 \text{ L}^2 \text{ atm mol}^{-2}$ (c) $1.5 \text{ L}^2 \text{ atm mol}^{-2}$ (d) $3.0 \text{ L}^2 \text{ atm mol}^{-2}$

Solution :

For one mole of a gas, the van der Waals equation is
$$\left(p + \frac{a}{V^2}\right)(V - b) = RT$$

When $b = 0$, we have
$$\left(p + \frac{a}{V^2}\right)V = RT \quad \text{i.e.} \quad pV = RT - \frac{a}{V}$$

The plot of pV versus $1/V$ will be a straight line with slope equal to $-a$. Hence

Slope of the given straight line is
$$\frac{(20.1 - 21.6) \text{ L atm mol}^{-1}}{(3.0 - 2.0) \text{ mol L}^{-1}} = -1.5 \text{ L}^2 \text{ atm mol}^{-2}$$

Equating this to $-a$, we get
$$a = 1.5 \text{ L}^2 \text{ atm mol}^{-2}$$

Therefore, the **choice (c)** is correct.

dilute solution containing 2.5 gm of nonvolatile

For a dilute solution containing 2.5 g of a non-volatile non-electrolyte solute in 100 g of water, the elevation of boiling point at 1 atm pressure is 2 °C. Assuming concentration of solute is much lower than the concentration of solvent, the vapour pressure of the solution (take $K_b = 0.76 \text{ K kg mol}^{-1}$) is

- (a) 724 mmHg (b) 740 mmHg (c) 736 mmHg (d) 718 mmHg

Solution :

From the expression $\Delta T_b = K_b m$, we find that the molality of solute (m) in the solution is

$$m = \frac{\Delta T_b}{K_b} = \frac{2 \text{ K}}{0.76 \text{ K kg mol}^{-1}} = \frac{2}{0.76} \text{ mol kg}^{-1}$$

Since $m = n_2/m_1$ (where n_2 is the amount of solute and m_1 is the mass of solvent expressed in kg), we get

$$n_2 = mm_1 = \left(\frac{2}{0.76} \text{ mol kg}^{-1}\right) (0.1 \text{ kg}) = \frac{2}{7.6} \text{ mol}$$

From the expression $-\Delta p = x_2 p_1^*$ of the relative lowering of vapour pressure of solvent (whose vapour pressure is 1 atm as the solute is non-volatile), we find that

$$x_2 = \frac{n_2}{n_1 + n_2} \approx \frac{n_2}{n_1} = \frac{n_2}{m_1/M_1} = \frac{(2/7.6) \text{ mol}}{(100 \text{ g}/18 \text{ g mol}^{-1})} = \frac{36}{76}$$

$$\text{Hence } -\Delta p = \left(\frac{36}{760}\right) (760 \text{ mmHg}) = 36 \text{ mmHg}$$

$$p = p_1^* + \Delta p = 760 \text{ mmHg} - 36 \text{ mmHg} = 724 \text{ mmHg}$$

Therefore, the **choice (a)** is correct.


shape of XeO_2F_2 molecule is seesaw

The shape of XeO_2F_2 molecule is


- (a) trigonal bipyramidal (b) square planar (c) tetrahedral (d) see-saw

Solution :

The number of valence electrons in XeO_2F_2 is $8 + 2 \times 6 + 2 \times 7 = 34$. These are distributed as shown in the following.



There are five pairs of electrons around Xe. To accommodate these, Xe undergoes dsp^3 hybridization leading to the trigonal bipyramidal arrangement of five pairs of electrons. The structure of XeO_2F_2 is



The shape of XeO_2F_2 is sea-saw.

Therefore, the **choice (d)** is correct.

Various increasing and Decreasing properties

Decreasing order of reactivity towards S_N2 displacement:

n-Butyl bromide, isobutyl bromide, *sec*-butyl bromide, *tert*-butyl bromide

Increasing order of reactivity towards S_N1 displacement:

1-bromopentane, 2-bromopentane, 2-bromo-2-methylbutane

Increasing reactivity sequence of S_N1 displacement of halogen: CH_3X , $1^\circ X$, $2^\circ X$, $3^\circ X$,

Increasing reactivity sequence of S_N2 displacement of halogen: CH_3X , $1^\circ X$, $2^\circ X$, $3^\circ X$.

Increasing reactivity of alcohols towards gaseous HBr:

2-butanol, 2-methyl-1-propanol, 2-methyl-2-propanol

Increasing order of reactivity towards E2 dehydrohalogenation of the bromides:

ethyl bromide, *n*-propyl bromide, isobutyl bromide, neopentyl bromide

Increasing stability of alkenes: $R_2C=CR_2$, $R_2C=CHR$, $R_2C=CH_2$, $RCH=CH_2$, $CH_2=CH_2$

Increasing dehydration of alcohol in the presence of H_2SO_4 :

ethyl alcohol, isopropyl alcohol, *tert*-butyl alcohol

Decreasing order of reactivity towards E2 dehydrohalogenation:


2-bromo-2-methylbutane, 1-bromopentane, 2-bromopentane, 3-bromopentane

Increasing order of dehydrohalogenation

1-bromo-3-methylbutane, 2-bromo-2-methylbutane, 3-bromo-2-methylbutane

calculate multiple bond energy of a C triple bond

Using the data provided, calculate the multiple bond energy of a $C\equiv C$ bond in C_2H_2 .




Take the bond energy of C—H bond equal to 350 kJ mol^{-1} .

- (a) 1165 kJ mol^{-1} (b) 837 kJ mol^{-1} (c) 865 kJ mol^{-1} (d) 815 kJ mol^{-1}

Solution:

Consider the following transformations.



According to Hess's law $\Delta H_1 = \Delta H_2 + \Delta H_3 - 2\epsilon_{C-H} - \epsilon_{C\equiv C}$

This gives

$$\epsilon_{C\equiv C} = \Delta H_2 + \Delta H_3 - \Delta H_1 - 2\epsilon_{C-H} = (1410 + 330 - 225 - 2 \times 350) \text{ kJ mol}^{-1} = 815 \text{ kJ mol}^{-1}$$

Therefore, the **choice (d)** is correct.

Some facts student must know

Abscisic acid (ABA) - A PLANT HORMONE once thought to be responsible for the shedding (abscission) of flowers and fruit and for the onset of dormancy in buds (hence its early name, dormin). The compound is associated with the closing of pores (stoma) in the leaves of plants deprived of water.

Absolute configuration - A particular molecular configuration of a CHIRAL molecule, as denoted by comparison with a reference molecule or by some sequence rule. There are two systems for expressing absolute configuration in common use: the D-L convention and the R-S convention.


Absorption

1. A process in which a gas is taken up by a liquid or solid, or in which a liquid is taken up by a solid. In absorption, the substance absorbed goes into the bulk of the material. Solids that absorb gases or liquids often have a porous structure. The absorption of gases in solids is sometimes called sorption. There is a distinction between absorption (in which one substance is assimilated into the bulk of another) and ADSORPTION (which involves attachment to the surface). Sometimes it is not obvious which process is occurring. For example, a porous solid, such as activated CHARCOAL may be said to absorb a large volume of gas, but the process may actually be adsorption on the high surface area of internal pores in the material.


2. The process in which electromagnetic radiation, particles, or sound waves lose energy in passing through a medium. Absorption involves conversion of one form of energy into another.

Accelerator - A substance that increases the rate of a chemical reaction. In this sense the term is synonymous with CATALYST. It is common to refer to catalysts as 'accelerators' in certain industrial applications. For example, accelerators are used in the VULCANIZATION of rubber and in the polymerization of adhesives. Also, in the production of composite materials using polyester resins a distinction is sometimes made between the catalyst (which initiates the polymerization reaction) and the accelerator (which is an additional substance making the catalyst more effective). The terms promoter and activator are used in a similar way.


Acceptor - The atom or group to which a pair of electrons is donated in forming a COORDINATE BOND.



Acenaphthene - ($C_{12}H_{10}$) A colorless crystalline derivative of naphthalene, used in producing some dyes.




Acetal - A type of compound formed by reaction of an alcohol with either an aldehyde or a ketone. The first step in formation of an acetal is the formation of an intermediate, known as a hemiacetal. For example, ethanal (acetaldehyde; CH_3CHO) reacts with ethanol ($\text{C}_2\text{H}_5\text{OH}$) as follows: $\text{CH}_3\text{CHO} + \text{C}_2\text{H}_5\text{OH} \rightleftharpoons \text{CH}(\text{OH})(\text{CH}_3)(\text{C}_2\text{H}_5\text{O})$ The **hemiacetal** has a central carbon atom (from the aldehyde) attached to a hydrogen, a hydroxyl group, a hydrocarbon group (methyl - CH_3), and an alkoxy group ($\text{C}_2\text{H}_5\text{O}$). If a ketone is used rather than an aldehyde, the resulting hemiacetal contains two hydrocarbon groups. For example, reaction of the ketone R^1COR^2 with the alcohol R^3OH is:
 $\text{R}^1\text{COR}^2 + \text{R}^3\text{OH} \rightleftharpoons \text{CR}^1\text{R}^2(\text{OH})(\text{OR}^3)$



The formation of a hemiacetal is an example of **NUCLEOPHILIC ADDITION** to the carbonyl group of the aldehyde or ketone. The first step is attack of the lone pair on the O of the alcohol on the (positively charged) C of the carbonyl group. This is catalyzed by both acids and bases. Acid catalysis occurs by protonation of the O on the carbonyl, making the C more negative and more susceptible to nucleophilic attack. In base catalysis the OH^- ions from the base affect the -OH group of the alcohol, making it a more effective nucleophile.

In general, hemiacetals exist only in solution and cannot be isolated because they easily decompose back to the component alcohol and aldehyde or ketone. However, some cyclic hemiacetals are more stable. For example, cyclic forms of **SUGAR** molecules are hemiacetals.

Further reaction of hemiacetals with another molecule of alcohol leads to a full acetal. For example:
 $\text{CH}(\text{OH})(\text{CH}_3)(\text{OC}_2\text{H}_5) + \text{C}_2\text{H}_5\text{OH} \rightleftharpoons \text{CH}(\text{CH}_3)(\text{OC}_2\text{H}_5)_2$




The overall reaction of an aldehyde or ketone with an alcohol to give an acetal can be written: $\text{R}^1\text{COR}^2 + \text{R}^3\text{OH} \rightleftharpoons \text{CR}^1\text{R}^2(\text{OR}^3)_2$

It is also possible to have 'mixed' acetals with the general formula $\text{CR}^1\text{R}^2(\text{OR}^3)(\text{OR}^4)$. Note that if the acetal is derived from an aldehyde, then R^1 and/or R^2

may be a hydrogen atom. The mechanism of formation of an acetal from a hemiacetal is acid catalyzed. It involves protonation of the -OH group of the hemiacetal followed by loss of water to form an oxonium ion, which is attacked by the alcohol molecule.

Heterolytic fission



Heterolysis

In this case we can see that one of the atoms carry a negative charge after bond cleavage indicating that it has both the electrons of the bond and the other has no electrons at all. Hence it is electron deficient thus positively charged. As the electrons are not divided equally after bond cleavage this is called *Heterolytic Fission*. In a case the C atom carries a positive charge it is called a carbocation and in the case it carries both the electrons of the broken bond and is negatively charged, it is called a Carbanion. Carbocation and Carbanions are the most important carbon intermediates in organic chemistry. Please note that both types of fissions are applicable to both homoatomic and heteroatomic bonds (bonds between two different atoms say C-N or C-O).


Free radicals

These are neutral intermediates, formed due to homolytic cleavage of a single bond. Some common bonds which cleave to give free radicals in organic chemistry are shown: C-O, C-Cl, C-Br, C-I, C-C, C-H. Carbon free radicals are mainly generated by:

- Photolysis (action of light) like acetone alpha cleavage
- Other radical initiator like allylic bromination by N-Bromosuccinimide (NBS)

There has been a certain degree of debate as to what the shape and geometry of a free radical is like. Revisiting the theory of hybridization, there can be two basic shapes of these radicals.


If the centre carbon atom of the radical is sp^3 hybridized (remember the one which was made of one s and three orbitals as in CH_4), the geometry will be tetrahedral.¹ But in the case of a radical there are only three groups attached to the sp^3 hybridized carbon atom so they will have a shape of what resembles a pyramid—it's a tetrahedron with its head cut off. So sp^3 hybridized radicals are pyramidal in shape. The single electron of the radical would then be housed in a sp^3 orbital. The other option is sp^2 hybridization. In that case the C atom is sp^2 hybridized, so as discussed previously the shape would be planar with the single electron in the unhybridized p-orbital with the three substituents having sp^2 hybridized bonds.



Two different geometries of free radicals. The single electrons are shown as black dots.

So to summarize free radicals:


- Formed under activation by light or use of additional compounds called Radical Initiators.
- They are very reactive, because they have an unpaired electron which wants to get paired up.
- They are either pyramidal or planar with the lone electron in their sp^3 or p orbitals respectively.
- Because of their high reactivity, they tend to be less selective. In simple terms it means that it sometimes difficult to predict what products are formed in reactions which involve free radicals and we actually get several products from a single reaction.



This reaction shows the formation of two products with the Br atom attached to different carbons.

Carbocations

Carbocations are formed from the heterolytic cleavage of a carbon-heteroatom (meaning a non carbon atom in general) bond where the other atom is more electronegative than carbon like a C-O, C-N, C-X (X can be Cl, Br, I, etc) bond. This is quite logical as after the cleavage if a carbocation is to be formed the two electrons of the bond must go to the other atom. And this is favoured if that other atom is electronegative. Carbocations can be made in difficult conditions by using so-called superacids, developed by George Olah (Nobel Prize, 1994), which helps stabilize these intermediates substantially to be analyzed. Formation of carbocations can be assisted by using cations like Ag^+ , with alkyl halides as substrates.



The precipitating out of the silver salt forces the equilibrium to shift towards the forwards reaction.

A Similar reaction is [Finkelstein reaction](#) is particularly useful for preparing iodoalkanes. The iodoalkanes are obtained by heating chloro or bromoalkanes with a concentrated solution of sodium iodide in dry acetone.

In Acetone after Reflux (Heating in enclosed container) for a long time [Refluxing is a process of heating a liquid in a flask provided with a condenser. The vapours of the liquid which rise up get condensed back into the flask.]

$\text{RX} + \text{NaI} \longrightarrow \text{RI} + \text{NaX}$ where $\text{X} = \text{Cl}, \text{Br}$, and R is an Alkyl group

Sodium chloride and sodium bromide being less soluble in acetone get precipitated from the solution and can be removed by filtration. This also prevents the backward reaction to occur according to Le Chatelier's principle. The reaction gives the best results with primary halides.

With iodide, the solvate $\text{NaI} \cdot 3(\text{CH}_3)_2\text{CO}$ can be formed, in which each Na^+ is coordinated by 6 acetone ligands via oxygen lone pairs, and the I^- ions fill in the gaps between these octahedral units, being surrounded by the methyl groups of the acetones. I^- has a larger ionic radius compared to Cl^- and Br^- and thus fills the gap completely, thereby minimizing empty space in the lattice and its distance to Na^+ , which is, however, still larger than in solvent-free sodium iodide. This larger distance between the ions decreases the lattice energy, which in turn makes the compound readily soluble in acetone. Cl^- and Br^- are too small to fit in these gaps, and as the cation-anion distance becomes too large, the only way to minimize the lattice energy is the formation of a solvent-free structure NaX ($X = \text{Cl}^-$, Br^-). When the lattice energy is larger than the energy released by solvation of the ions, namely that of Na^+ by complexation with acetone, the compound becomes insoluble and precipitates from the solution.

The positively charged carbon atom in carbocations is sp^2 hybridized, which means it's planar as we know by now. The three substituents of the carbocation lie in a plane leaving the unhybridized empty p orbital perpendicular to them.

These intermediates react with species which are electron rich (quite obvious) and being charged are stabilized in polar solvents. (Just as Na^+ is soluble and stable in polar water). Carbocations are important intermediates in most mechanisms along with carbanions as we shall see later.

To summarize carbocations:

- Formed due to heterolysis of a C-X bond (where X is more electronegative) and thus has a positive charge.
- Planar in shape (sp^2 hybridized carbon), with empty p orbital perpendicular to the plane of the molecule.
- Reactive towards electron rich species.

Carbanions

These are intermediates also formed as a result of heterolysis, but here the electron pair from the bond is kept by the carbon atom. From what we saw earlier the more electronegative atom keeps the electrons, so in this case carbon must be the more electronegative of the two atoms making up the bond. Now there are only a few atoms (non-metals; metals are not usually part of organic chemistry) which are less electronegative, so the most common bond cleavage which yields carbanions is the C-H bond. The ease of breaking this bond and creating a carbanion is also a measure of the compound's acidity, because a H^+ is also generated with the carbanion, which makes the molecule an acid in the Bronsted sense.

Carbanions have three groups attached to each other and a lone pair of electrons which gives it its negative charge (similar to the ammonia molecule where the central N has 3 Hs and a lone pair of electrons). So its geometry is pyramidal (tetrahedral but since there is no fourth group again it's like a tetrahedral with head cut off) and the carbon atom is sp^3 hybridized.

Carbanions are also stable in polar solution (electrostatic stabilization).


To summarize carbanions:

- Formed due to heterolysis of a C-X bond (where X is less electronegative) and thus has a negative charge.
- Pyramidal in shape (sp^3 hybridized) with the excess electrons placed in one sp^3 hybrid orbital.
- Reactive towards positively charged (electron deficient species).


Stability of intermediates

Most organic reactions take place via formation of intermediates. So the study of different intermediates would help us predict the course of the reaction and the main aspect to look at would be their stability. No organic mechanism has been conclusively 'PROVEN', all the mechanism we see are the most plausible ones derived from many experiments, a major component of which is isolating and studying the intermediates. It is difficult to say that a certain mechanism is absolutely correct, but it is quite simple to point out an incorrect mechanism. One of the ways a chemist would confirm an incorrect mechanism is if it involves a very unstable intermediate. The good thing about this is that with a few empirical rules and principles in mind, it is quite simple to assign relative stability of intermediates like radicals, carbocations and carbanions. And what is even better is that we have already discussed these principles.

For carbocations and free radicals (both electron poor species), any group which donates electron density to the carbon centre would stabilize it and inversely electron withdrawing groups would increase electron deficiency on the carbon centre leading to destabilization. (Remember charge is not desirable, the most stable species are usually neutral). So following the same logic the effect should just be opposite in the case of carbanions as they are electron rich (negatively charged) instead of being electron deficient like the above two. So groups which pull away electrons from the charged carbon atom would have a stabilizing effect whereas electron donation would destabilize the intermediate as it loads more negative charge on an already negatively charged atom.





The name of a complex radical is considered to start with the first letter of its complete name.





dimethylpentyl (as a complete single substituent) is alphabetized under "d", thus 7-(1,2-Dimethylpentyl)-5-ethyltridecane


In cases where names of complex radicals are composed of identical words, preference for citation is given to that radical which contains the lowest locant at the first cited point of difference in the radical.




If two or more side chains are in equivalent positions the one to be assigned the lower number is that cited first in the name.





The lowestsum rule.




2,3,5-Trimethyl-4-propylheptane (not 3,5,6-Trimethyl-4-propylheptane)



4-Isobutyl-2,5-dimethylheptane (not 4-Isobutyl-2, 6-dimethylheptane)




CLASSIFICATION OF ORGANIC COMPOUNDS



Alicyclic or closed chain or ring compounds


Alicyclic (aliphatic cyclic) compounds contain carbon atoms joined in the form of a ring (homocyclic). Sometimes atoms other than carbon are also present in the ring (heterocyclic).

Some examples are




Nomenclature of Substituted Benzene Compounds

For IUPAC nomenclature of substituted benzene compounds, the substituent is placed as prefix to the word *benzene* as shown in the following examples.




How Isomers are made ?

(a) *Normal to branched chain in alkane* *n*-Alkane is heated with AlCl_3 at 570 K to give branched chain




(b) *Shift of double bond*

In the presence of catalyst like $\text{Al}_2(\text{SO}_4)_3$, alkene undergo isomerization at high temperature (770 – 970 K).




(c) *Shift of triple bond*




Heating with ethanolic potassium hydroxide shifts the triple bond towards the centre of the chain whereas heating with sodamide in an inert solvent shifts the bond towards the end of chain.

The choice of principal functional group is made on the basis of order of preference. **The order of decreasing priority for some functional groups is: $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{COOR}$ ($\text{R}=\text{alkyl group}$), COCl , $-\text{CONH}_2$, $-\text{CN}$, $-\text{HC}=\text{O}$, $>\text{C}=\text{O}$, $-\text{OH}$, $-\text{NH}_2$, $>\text{C}=\text{C}<$, $-\text{C}\equiv\text{C}-$.**



Epimers



Anomers









Epimer Anomer comparison



Epimers are monosaccharides that differ in chirality at only one carbon. In the straight-chain format, epimers will have H and OH-substituents switched at one backbone carbon, but not at any others.

Anomers are special epimers; in cyclic forms of one single monosaccharide, anomers differ in chirality at the anomeric (hemiacetal) carbon only. In the straight-chain format, anomers will have the same configuration.

Various Methods of Reduction used in Organic Chemistry







(j) Rosenmund's reduction




Various methods of Oxidation

Oxidation




Synthesis of *p*-bromonitrobenzene in 2 steps

a synthesis of *p*-bromonitrobenzene from benzene in two steps.




Secondary alcohol to tertiary alcohol




Prepare *m*-bromiodobenzene from benzene


prepare *m*-bromiodobenzene from benzene (in not more than 5-7 steps)?



Alternatively, we may proceed as follows:




Secondary alcohol to Primary alcohol




To convert Ethylbenzene to Benzene and Ethylbenzene to phenylpropionic acid

Show the steps to carry out the following transformations.


(i) Ethylbenzene → benzene



(ii) Ethylbenzene → 2-phenylpropionic acid.




Alcohol to alcohol with one carbon more




Carry out the transformations 1-Butyne to ketone


Carry out the following transformation in not more than three steps.



Alcohol to alcohol with one carbon less





Simple alcohol to complicated alcohol



Conversion of Aniline to Benzylamine

How would you bring about the following conversion (in 3 steps)?

Aniline \rightarrow Benzylamine



Methanoic acid



Ethanoic acid

Propanoic acid

(i) Carbon-oxygen bond lengths in formic acid are different but are the same in sodium formate. Justify.

(ii) Phenate ion has more number of contributing structures than benzoate ion: but still benzoic acid is a stronger acid. Explain.


Solution. (i) In sodium formate, the contributing structures for the anion are equivalent while these are not the same in formic acid.



Thus, carbon-oxygen bond length in formate ion is the same for both the bonds while these are different in formic acid.


(ii) In phenate ion, the negative charge is dispersed only one electronegative oxygen atoms while there are two available oxygen atoms in benzoate ion to disperse the negative charge.

Amine to amine containing one carbon more




Synthesis of 4-methoxyphenol from bromobenzene

How would you synthesise 4-methoxyphenol from bromobenzene is NOT more than five steps? State clearly the reagents used in each step and show the structure of the intermediate compounds in your synthetic scheme.



The introduction of $-\text{NO}_2$ group at *para* position in bromobenzene facilitates the nucleophilic replacement of $-\text{Br}$ by $-\text{OCH}_3$.



The negative charge is strongly stabilised by resonance involving *para*-nitro group.

Write chemical tests to distinguish

between:


Solution. Formic acid $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{C}-\text{OH} \end{array} \right)$ contains

both an aldehyde $\left(\begin{array}{c} \text{H} \\ | \\ -\text{C}=\text{O} \end{array} \right)$ as well as carboxyl

group $\left(\begin{array}{c} -C-OH \\ || \\ O \end{array} \right)$ but acetic acid contain only a


carboxyl group. Formic acid behaves as reducing agent whereas acetic acid does not.

(a) Formic acid reduces Tollen's reagent to metallic silver but acetic acid does not.




No silver mirror is formed with acetic acid.

(b) Formic acid reduced Fehling solution to red ppt. of Cu_2O but acetic acid does not.



Amine to amine containing one carbon less





Convert Benzoic acid to meta fluorobenzoic acid (2-fluorobenzoic acid)

Convert




metafluoro benzoic acid






Aldehyde to aldehyde containing one carbon more




Convert Nitrobenzene to meta hydroxynitrobenzene (or 4-nitrophenol)





The given conversion can be carried out as follows.

Nitrobenzene resonance structure



4 nitrophenol or metahydroxynitrobenzene









Many conversions or preparations


Write down the reactions involved in the preparation of the following using the reagents indicated against each in paranthesis

- (i) Ethylbenzene from benzene [$\text{C}_2\text{H}_5\text{OH}$, PCl_5 , anhydrous AlCl_3]
 (ii) Propanoic anhydride from propanol [$\text{AgNO}_3/\text{NH}_4\text{OH}$, P_2O_5]
 (iii) Acetoxime from acetaldehyde [$\text{K}_2\text{Cr}_2\text{O}_7/\text{H}^+$, $\text{Ca}(\text{OH})_2$ and $\text{NH}_2\text{OH} \cdot \text{HCl}$]




Benzene to p-toluic acid

Benzene to p-toluic acid




Benzene to m-chlorobenzoic acid

Benzene to m-chlorobenzoic acid




Benzene to *p*-chlorobenzoic acid

Benzene to *p*-chlorobenzoic acid




Acetic acid to Malonic acid

Acetic acid to malonic acid




Ethyne to propanoic acid

Ethyne to propanoic acid




Isopropyl alcohol to β -hydroxybutyric acid

Isopropyl alcohol to β -hydroxybutyric acid





Acetic acid to Propanoic acid

Acetic acid to propanoic acid




Ethanoic acid to a mixture of methanoic acid and diphenyl ketone

Ethanoic acid to a mixture of methanoic acid and diphenyl ketone




Propanoic acid to acetic acid

Propanoic acid to acetic acid




Aniline to 1, 2, 3-tribromobenzene




Ethanal to 2-hydroxy-3-butenic acid

Ethanal to 2-hydroxy-3-butenic acid




Benzene to m-bromophenol

Benzene to m-bromophenol




Ethyl alcohol to Vinyl alcohol

Ethyl alcohol to vinyl acetate




Methylamine to ethylamine


Methylamine to ethylamine




Benzene to *m*-bromotoluene




Benzaldehyde to cyanobenzene



Aniline to Chlorobenzene



Ethylamine to Methylamine




Benzene to *m*-dibromobenzene

Benzene to *m*-dibromobenzene




Aniline to *p*-nitroaniline

Aniline to *p*-nitroaniline





Toluene to *m*-nitrotoluene

Toluene to *m*-nitrotoluene




Convert Benzene + Alcohol to α -methylphenylacetic acid

Benzene + alcohol to α -methylphenylacetic acid




Toluene to 3,5-dibromotoluene

Toluene to 3,5-dibromotoluene




Benzene to p-chloronitrobenzene

Benzene to *p*-chloronitrobenzene




Toluene to 2,6-dibromotoluene




Benzene to 3,4-dibromonitrobenzene

Benzene to 3,4-dibromonitrobenzene




Benzene to *m*-chloronitrobenzene

Benzene to *m*-chloronitrobenzene




Nitrobenzene to benzamide

Nitrobenzene to benzamide





Benzene to 4-nitrobenzaldehyde

Benzene to 4-nitrobenzaldehyde



Benzene to 4-amino-2-bromotoluene



Benzene to 4-amino-2-bromotoluene




2,4-Dinitrophenol (DNP)

Convert a mixture of *cis*-2-pentene and *trans*-2-pentene into *cis*-2-pentene or *trans*-2-pentene


A mixture of *cis*-2-pentene and *trans*-2-pentene into *cis*-2-pentene or *trans*-2-pentene




Benzene to *p*-chlorobenzylalcohol





Convert Propanol to α -hydroxypropanoic acid



4-chloropentane-2-one




3-Bromo-4-methylpentanoic acid



Vanillin




Salicylaldehyde




Cinnamaldehyde

p,p'- dihydroxybenzophenone




Hex-2-ene-4-ynoic acid




Heptan-2-one




Benzene to Mandelic Acid



4-Bromo-2-methylhexane




Heptanal





1-(4-Methoxyphenyl)-3-phenylprop-2-en-1-one

1-(4-Methoxyphenyl)-3-phenylprop-2-en-1-one




4-phenylprop-3-ene-2-one




CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

3-phenylprop-2-ene-1-al




3-phenyl-prop-2-ene-1-ol



cyclopentancarbaldehyde




Benzophenone also diphenylmethanone




CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



Tollen's Reagent





An aldehyde



Free silver





Benzene to 2,4-dinitrobenzaldehyde



Acetaldehyde to Acetylene

Acetaldehyde to acetylene






Identify the compounds


compound contains C, H and O unreactive towards Na (Sodium)

A compound (X) containing only C, H and O is unreactive towards sodium. It does not add bromine. It also does not react with Schiff's reagent. On refluxing with an excess of hydroiodic acid, X yields only one organic product Y. On hydrolysis Y yields a new compound Z which can be converted to Y by reaction with red phosphorus and iodine. The compound Z on oxidation with potassium permanganate gives a carboxylic acid. The equivalent weight of this acid is 60. What are the compounds X, Y and Z? Write chemical equations leading to the conversion of X to Y. (1981)





Solution Since the compound is unreactive towards sodium, it is neither alcohol (including phenol) nor carboxylic acid. It is also not an aldehyde. It may be a symmetrical ether because on refluxing with excess of HI, it gives only one product. Thus, we have



Now the equivalent weight of R'COOH is 60. From this, it follows that R' is —CH₃ group and hence R is —CH₂CH₃ group. Thus, the compounds X, Y and Z are



Steric Effects and the Relative Reactivity of Carboxylic Acid Derivatives




Identify the compound


A hydrocarbon reacts with Bromine to give dibromo compound

One mole of a hydrocarbon A reacts with one mole of bromine giving a dibromo compound $C_5H_{10}Br_2$. Substance A on treatment with cold, dilute alkaline potassium permanganate solution forms a compound $C_5H_{12}O_2$. On ozonolysis, A gives equimolar quantities of propanone and ethanal. Deduce the structural formula of A.

Solution From the products of ozonolysis, we conclude that



The compound A contains a double bond. This is also confirmed from the fact that the compound A reacts with one mole of bromine. Thus, the given reactions are




Identify the compound


An alkene A on ozonolysis yields acetone and an aldehyde

An alkene A on ozonolysis yields acetone and an aldehyde. The aldehyde is easily oxidized to an acid B. When B is treated with bromine in presence of phosphorus, it yields a compound C which on hydrolysis gives a hydroxy acid D. This acid can also be obtained from acetone by the reaction with hydrogen cyanide followed by hydrolysis. Identify the compounds A, B, C and D.


Solution The structure of compound D is easily deduced from the fact that it is obtained by the combination of acetone with hydrogen cyanide followed by hydrolysis.




The conversion B to C and then C to D indicates that the reaction B to C is Hell-Volhard-Zelinsky reaction. Thus, we have





Hence, we conclude that the aldehyde from which compound B is obtained by oxidation has the structure



Now, the ozonolysis of the compound A gives acetone and 2-methyl-propanal, i.e.





Hence, the compound A, B, C and D are



2, 4-dimethylpent-2-ene

2-methylpropanoic acid



2-bromo-2-methylpropanoic acid



2-hydroxy-2-methylpropanoic acid

Identify the compound

An aromatic compound contains 69.4% carbon, 5.8% hydrogen

An aromatic compound contains 69.4% carbon and 5.8% hydrogen. A sample of 0.303 g of this compound was analysed for nitrogen by Kjeldahl's method. The ammonia evolved was absorbed in 50 mL of 0.05 M sulphuric acid. The excess acid required 25 mL of 0.1 M sodium hydroxide for neutralization. Determine the molecular formula of the compound if its molar mass is 121 g mol^{-1} . Draw two possible structures for this compound.

Solution The reactions involved are



$$\text{Amount of sodium hydroxide to neutralize excess of sulphuric acid} = (25 \text{ mL}) \left(\frac{0.1 \text{ mol}}{1000 \text{ mL}} \right) = \frac{1}{400} \text{ mol}$$

$$\text{Amount of sulphuric acid neutralized with sodium hydroxide} = \frac{1}{2} \left(\frac{1}{400} \text{ mol} \right) = \frac{1}{800} \text{ mol}$$

$$\text{Amount of sulphuric acid taken to start with} = (50 \text{ mL}) \left(\frac{0.05 \text{ mol}}{1000 \text{ mL}} \right) = \frac{1}{400} \text{ mol}$$

$$\text{Amount of sulphuric acid neutralized by ammonia} = \left(\frac{1}{400} - \frac{1}{800} \right) \text{ mol} = \frac{1}{800} \text{ mol}$$

$$\text{Amount of ammonia evolved} = 2 \left(\frac{1}{800} \text{ mol} \right) = \frac{1}{400} \text{ mol}$$

$$\text{Mass of nitrogen in the evolved ammonia} = \left(\frac{1}{400} \text{ mol} \right) (14 \text{ g mol}^{-1}) = \frac{7}{200} \text{ g}$$

$$\text{Per cent of nitrogen in the given aromatic compound} = \frac{(7/200) \text{ g}}{0.303 \text{ g}} \times 100 = 11.6$$

$$\text{Per cent of oxygen in the given aromatic compound} = 100 - (69.4 + 5.8 + 11.6) = 13.2$$

The ratios of atoms present in the given compound are

$$\text{C} : \text{H} : \text{N} : \text{O} \quad :: \quad \frac{69.4}{12} : \frac{5.8}{1} : \frac{11.6}{14} : \frac{13.2}{16} \quad :: \quad 5.8 : 5.8 : 0.83 : 0.83 \quad :: \quad 7 : 7 : 1 : 1$$

Hence, for the given compound, we have


$$\text{Empirical formula} = \text{C}_7\text{H}_7\text{NO}$$

$$\text{Empirical molar mass} = 121 \text{ g mol}^{-1}$$

It is given that 121 g mol^{-1} is also the molar mass. Hence,

$$\text{Molecular formula} = \text{C}_7\text{H}_7\text{NO}$$

Since, the given compound is aromatic, the two possible structures are



Identify the compound

An Organic compound $\text{C}_x\text{H}_2y\text{O}_y$ was burnt with twice the amount of Oxygen

An organic compound $\text{C}_x\text{H}_2y\text{O}_y$ was burnt with twice the amount of oxygen needed for complete combustion to CO_2 and H_2O . The hot gases, when cooled to 0°C and 1 atm pressure, measured 2.24 L. The water collected during cooling weighed 0.9 g. The vapour pressure of pure water at 20°C is 17.5 mmHg and is lowered by 0.104 mmHg when 50 g of the organic compound is dissolved in 1000 g of water. Give the molecular formula of the organic compound.

Solution The combustion reaction is $\text{C}_x\text{H}_2y\text{O}_y + x \text{O}_2 \rightarrow x \text{CO}_2 + y \text{H}_2\text{O}$

To start with, the amount of O_2 taken is $2x$. Hence, after the combustion reaction, we will be left with the following amounts.

$$\text{Amount of oxygen left unreacted} = x; \quad \text{Amount of carbon dioxide} = x; \quad \text{Amount of water} = y$$

When this mixture is cooled to 0°C and 1 atm, we will be left with oxygen and carbon dioxide. Hence, the amount $2x$ occupies the given volume of 2.24 L at STP. Hence,

$$\text{Amount } x = \frac{(2.24/2) \text{ L}}{22.4 \text{ L mol}^{-1}} = 0.05 \text{ mol}$$

$$\text{Now,} \quad \text{Mass of water collected} = 0.9 \text{ g} \quad \text{Amount of water collected, } y = \frac{0.9 \text{ g}}{18 \text{ g mol}^{-1}} = 0.05 \text{ mol}$$

Thus, the empirical formula of the compound is $\text{C}_{0.05}\text{H}_{2 \times 0.05}\text{O}_{0.05}$, i.e. CH_2O . Now, according to Raoult's law

$$-\frac{\Delta p}{p^*} = x_2 \quad \text{i.e.} \quad \frac{0.104 \text{ mmHg}}{17.5 \text{ mmHg}} = \frac{(50 \text{ g}/M)}{(50 \text{ g}/M) + (1000 \text{ g}/18 \text{ g mol}^{-1})}$$

Identify the compound

Certain hydrocarbon was found to contain 85.7% carbon

A certain hydrocarbon A was found to contain 85.7 per cent carbon and 14.3 per cent hydrogen. This compound consumes 1 molar equivalent of hydrogen to give a saturated hydrocarbon B. 1.0 g of hydrocarbon A just decolourized 38.05 g of a 5 per cent solution (by mass) of Br₂ in CCl₄. Compound A, on oxidation with concentrated KMnO₄, gave compound C (molecular formula C₄H₈O) and compound C could easily be prepared by the action of acidic aqueous mercuric sulphate on 2-butyne. Determine the molecular formula of A and deduce the structures A, B and C.

Solution The ratio of atoms in the compound A is $C : H :: \frac{85.7}{12} : \frac{14.3}{1} :: 7.14 : 14.3 :: 1 : 2$

Thus, Empirical formula of A is CH₂.

Since the compound A consumes 1 mol of hydrogen, the molecule of A contains only one carbon-carbon double bond. From the data on the absorption of bromine, we can calculate the molar mass of A as shown in the following.


$$\text{Mass of bromine absorbed by 1.0 g of hydrocarbon} = \frac{5}{100} \times 38.05 \text{ g}$$

$$\text{Mass of hydrocarbon absorbing 160 g (= 1 mol) of Br}_2 = \frac{1.0}{(5 \times 38.05/100)} \times 160 \text{ g} = 84.1 \text{ g.}$$


Hence, Molar mass of A is 84.1 g mol⁻¹.

The number of repeating CH₂ group in one molecule of A will be 6 (= 84.1/14). Hence,

Molecular formula of A is C₆H₁₂. Now, it is given that



The compound C is obtained by the hydration of 2-butyne. Hence, its structure obtained from the reaction is




Identify the compound

A liquid X having a molecular formula C₆H₁₂O₂ is hydrolyzed with water


A liquid X, having a molecular formula C₆H₁₂O₂ is hydrolysed with water in the presence of an acid to give a carboxylic acid, Y, and an alcohol Z. Oxidation of Z with chromic acid gives Y. What are the structures of X, Y and Z?

Solution Since the hydrolysis of X gives a carboxylic acid (Y) and an alcohol (Z), the compound X must be an ester. Let it be RCOOR'. The compounds Y and Z will be RCOOH and R'OH, respectively. Since the oxidation of Z gives Y, we will have



Hence, the given compound X may be written as RCOOCH₂R. From this it follows that

$$2\text{R} = \text{C}_6\text{H}_{12}\text{O}_2 - \text{C}_2\text{H}_4\text{O} = \text{C}_4\text{H}_8 \quad \text{or} \quad \text{R} = \text{C}_2\text{H}_5$$



Identify the compound

An unknown compound of carbon, hydrogen, and oxygen contains 69.77% carbon and 11.63% hydrogen and has a molar mass of 86 g mol^{-1} . It does not reduce Fehling's solution, but forms a bisulphite addition compound and gives a positive iodoform test. What are the possible structures?

Solution The given compound is a ketone as it does not reduce Fehling's solution, but forms a bisulphite addition compound. It will also contain CH_3CO group as it shows positive iodoform test.


The ratios of atoms in the compound are

$$\text{C} : \text{O} : \text{H} :: \frac{69.77}{12} : \frac{18.60}{16} : \frac{11.63}{1} :: 5.81 : 1.16 : 11.63 :: 5 : 1 : 10$$

Hence Empirical formula is $\text{C}_5\text{H}_{10}\text{O}$.

Thus, Molecular formula is $\text{C}_5\text{H}_{10}\text{O}$.

The structure of the compound will be




Identify the compound

An organic compound A on treatment with acetic acid in the presence of Sulphuric acid


An organic compound A on treatment with acetic acid in the presence of sulphuric acid produces an ester B. The compound A on mild oxidation gives C. The compound C with 50% potassium hydroxide followed by acidification with dilute hydrochloric acid generates A and D. The compound D with phosphorus pentachloride followed by reaction with ammonia gives E. The compound E on dehydration produces hydrocyanic acid. Identify the compounds A, B, C, D and E.

Solution The given reactions are




The reaction $\text{C} \xrightarrow{50\% \text{ KOH}} \text{A} + \text{D}$ is a Cannizzaro reaction and thus the compound C is an aldehyde with no α -hydrogen, C must be HCHO. From this it follows that A must be CH_3OH as its mild oxidation gives HCHO.

With this fact, we can write the given reactions as shown below.



Hence



A : CH_3OH methanol






B : $\text{CH}_3\text{COOCH}_3$ methylethionate

C : HCHO methanal

D : HCOOH methanoic acid

E : HCONH_2 methanamide

Iodoform Test



Identify the compound

An organic compound X, on analysis gives 24.24 per cent carbon and 4.04 per cent hydrogen. Further, sodium extract of 1.0 g of X gives 2.90 g of silver chloride with acidified silver nitrate solution. The compound X may be represented by two isomeric structures, Y and Z. The compound Y on treatment with aqueous potassium hydroxide solution gives a dihydroxy compound, while Z on similar treatment gives ethanal. Find out the molecular formula of X and give the structures of Y and Z.

Solution Mass per cent of chlorine in the organic compound X

$$= \frac{M_{\text{Cl}}}{M_{\text{AgCl}}} \frac{m_{\text{AgCl}}}{m_{\text{compound}}} \times 100 = \frac{35.5}{143.5} \times \frac{2.90}{1.0} \times 100 = 71.21$$

The ratios of atoms in the molecule of X are

$$\text{C} : \text{H} : \text{Cl} :: \frac{24.24}{12} : \frac{4.04}{1} : \frac{71.72}{35.5} :: 2 : 4 : 2 :: 1 : 2 : 1$$



Empirical formula of X is CH_2Cl

Since the isomer Y of the compound gives a dihydroxy compound on treating with aqueous KOH, it follows that the compound must contain two Cl atoms. Hence,

Molecular formula of X is $\text{C}_2\text{H}_4\text{Cl}_2$.

Its two isomers are CH_3CHCl_2 and $\text{ClCH}_2\text{CH}_2\text{Cl}$.
(Z) (Y)

The reaction are



Match the entries given on the left with those given on the right.

- | | |
|---|--------------------------------|
| (a) Cyclopropane, chlorine and light | (p) Electrophilic substitution |
| (b) Propanone and sodium bisulphite | (q) Homologous pair |
| (c) Ethanal and Methanal | (r) Homolytic addition |
| (d) Benzene, nitric acid and sulphuric acid | (s) Free radical substitution |
| (e) Propene, hydrogen bromide and a peroxide catalyst | (t) nucleophilic addition |

Ans :

- (a) – (s); (b) – (t); (c) – (q); (d) – (p); (e) – (r)

<https://zookeepersblog.wordpress.com/science-tuition-chemistry-physics-mathematics-for-iit-jee-aieee-std-11-12-pu-isc-cbse/>

Match the entries given on the left with those given on the right.

- | | |
|--|-----------------------------|
| (a) Pyrolysis of alkanes | (p) Elimination reaction |
| (b) Benzene + Chloroethane
(+ anhydrous AlCl_3) | (q) Saponification |
| (c) $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{NaOH}$ | (r) Wurtz reaction |
| (d) Preparation of alkanes | (s) Friedel-craft reaction |
| (e) Phenol + CHCl_3 (NaOH) | (t) Reimer-Tiemann reaction |
| (f) $\text{C}_2\text{H}_5\text{Br} + \text{alcKOH}$ | (u) Cracking |


Ans :

- (a) – (u); (b) – (s); (c) – (q); (d) – (r); (e) – (t);
(f) – (p)

Identify the compound

A mixture of two aromatic compounds A and B was separated by dissolving in chloroform followed by extraction with aqueous KOH solution. The organic layer containing compound A, when heated with alcoholic solution of KOH produced a compound C ($\text{C}_7\text{H}_5\text{N}$) associated with an unpleasant odour. The alkaline aqueous layer on the other hand, when heated with chloroform and then acidified gave a mixture of two isomeric compounds D and E of molecular formula $\text{C}_7\text{H}_6\text{O}_2$. Identify the compounds A, B, C, D, and E and write their structures.


Solution We have



The compound C is an isocyanide which is produced from a primary amine. Hence, the structures of C and A are




The compound B exhibits Reimer-Tiemann reaction. Thus, the compound B must be phenol. The reaction is




Rearrangement by migration of Bromine

Rearrangement, by migration of Br, of initially formed 1° radical into more stable 2° or 3° radical.

Chlorination of either *n*-propyl or isopropyl bromide gives 1-bromo-2-chloropropane.



Chlorination of either isobutyl or *tert*-butyl bromide gives 1-bromo-2-chloro-2-methylpropane.



Identify the Compound

An organic compound containing C, H and O exists in two isomeric forms A and B. A mass of 0.108 g of one of the isomers gives on combustion 0.308 g of CO₂ and 0.072 g of H₂O. A is insoluble in NaOH and NaHCO₃ while B is soluble in NaOH. A reacts with concentrated HI to give compounds C and D. C can be separated from D by the ethanolic AgNO₃ solution and D is soluble in NaOH. B reacts readily with bromine to give compound E of molecular formula, C₇H₅OBr₃. Identify A, B, C, D and E with justification and give their structures.

Solution We have

$$\text{Per cent of carbon in the compound} = \frac{M_C}{M_{\text{CO}_2}} \frac{m_{\text{CO}_2}}{m_{\text{compound}}} \times 100 = \left(\frac{12}{44}\right) \left(\frac{0.308}{0.108}\right) (100) = 77.78$$

$$\text{Per cent of hydrogen in the compound} = \frac{2M_H}{M_{\text{H}_2\text{O}}} \frac{m_{\text{H}_2\text{O}}}{m_{\text{compound}}} \times 100 = \left(\frac{2}{18}\right) \left(\frac{0.072}{0.108}\right) (100) = 7.41$$


$$\text{Per cent of oxygen in the compound} = 100 - (77.78 + 7.41) = 14.81.$$

The ratios of atoms in the compound are


$$\text{C} : \text{H} : \text{O} :: \frac{77.78}{12} : \frac{7.41}{1} : \frac{14.81}{16} :: 6.48 : 7.41 : 0.926 :: 7 : 8 : 1$$

Hence, Empirical formula of the compound is C₇H₈O.

Since the isomer B on reacting with bromine water gives compound E (C₇H₅OBr₃), the molecular formula of A and B will be the same as the empirical formula derived above, since both contain the same number of carbon atoms. As E is obtained from B by the substitution of hydrogen with bromine and since there is high carbon content in B, the compounds A and B must be aromatic. Now, since compound A is insoluble in NaOH and NaHCO₃ and compound B is soluble in NaOH, it may be concluded that B is a phenolic compound and A is an ether. Hence, the structures of A and B are




(A)
anisole




(B)
m-cresol


The bromination of B gives



The reaction of compound A with HI is



The compound C can be separated from D by use of ethanolic AgNO₃ solution as it is soluble in it whereas D will remain insoluble. The compound D will be soluble in NaOH as it is phenol. Hence, the structures of A, B, C, D and E are



Match the entries given on the left with those given on the right.

- | | |
|----------------------------|-------------------------|
| (a) Decarboxylation | (p) Addition reaction |
| (b) Ozonolysis | (q) Sodalime |
| (c) Williamson's synthesis | (r) Structure of alkene |
| (d) Dichloroethylene | (s) Ether |

Ans :


(a) – (q); (b) – (r); (c) – (s); (d) – (p)

Identify the compound

Basic volatile Nitrogen compound

A basic volatile nitrogen compound gave a foul smelling gas when treated with chloroform and alcoholic potash. A 0.295 g sample of the substance, dissolved in aqueous HCl, and treated with NaNO₂ solution at 0 °C liberated a colourless, odourless gas whose volume corresponded to 112 mL at STP. After the evolution of the gas was complete, the aqueous solution was distilled to give an organic liquid which did not contain nitrogen and which on warming with alkali and iodine gave a yellow precipitate. Identify the original substance. Assume that it contains one N atom per molecule.

Solution Since the compound gives a foul smelling gas on treating with CHCl₃ and alcoholic KOH, the compound must be a primary amine.



Since the compound on treating with NaNO₂/HCl at 0 °C produces a colourless gas, the compound must be an aliphatic primary amine.



Thus, the gas produced is nitrogen.

$$\text{Amount of gas liberated} = \frac{112 \text{ mL}}{22400 \text{ mL mol}^{-1}} = \frac{1}{200} \text{ mol}$$

From the above equation, it is obvious that


$$\text{Amount of compound RNH}_2 = \frac{1}{200} \text{ mol}$$

If M is the molar mass of RNH_2 , then

$$\frac{2.295 \text{ g}}{M} = \frac{1}{200} \text{ mol} \quad \text{or} \quad M = 0.295 \times 200 \text{ g mol}^{-1} = 59 \text{ g mol}^{-1}$$

Thus, the molar mass of alkyl group R is $(59 - 16) \text{ g mol}^{-1}$, i.e. 43 g mol^{-1} . Hence, R must be C_3H_7 .

From Eq. (2), it is obvious that the liquid obtained after distillation is ROH . Since this gives yellow precipitate with alkali and iodine (iodoform test), it must contain $\text{CH}_3-\overset{|}{\underset{|}{\text{C}}}-\text{OH}$ group. Hence, it is concluded that ROH is



Match the entries given on the left with those given on the right.

- | | |
|-----------------------------|----------------------|
| (a) Lucas test | (p) Phenol |
| (b) Neutral FeCl_3 | (q) Glucose |
| (c) Dye test | (r) Tertiary alcohol |
| (d) Tollens test | (s) Aniline |


Ans :

(a) - (r); (b) - (p); (c) - (s); (d) - (q)

Identify the compound


An organic compound A, C_8H_6 , on treatment with dilute sulphuric acid containing mercuric sulphate gives a compound B, which can also be obtained from a reaction of benzene with an acid chloride in the presence of anhydrous aluminium chloride. The compound B, when treated with iodine in aqueous KOH , yields C and a yellow compound D. Identify A, B, C and D with justification. Show how B is formed from A.

Solution The given reactions are



The reaction of B with I_2 in KOH is iodoform reaction. The compound D is iodoform, CHI_3 . The compound B must contain $-\text{COCH}_3$ group so as to exhibit iodoform reaction. Since D is obtained from benzene by Friedel-Crafts reaction, it is an aromatic ketone ($\text{C}_6\text{H}_5\text{COCH}_3$). The compound C must be an acid.


The compound A may be represented as $\text{C}_6\text{H}_5\text{C}_2\text{H}$. Since it gives $\text{C}_6\text{H}_5\text{COCH}_3$ on treating with dilute H_2SO_4 and HgSO_4 , it must contain a triple bond ($-\text{C}\equiv\text{CH}$) in the side chain. Hence, the given reactions may be represented as




Identify the compound

An organic compound E (C_5H_8) on hydrogenation gives compound F (C_5H_{12}). Compound E on ozonolysis gives formaldehyde and 2-keto-propanal. Deduce the structure of compound E.

Solution Since there are four hydrogen atoms added in the reaction




the compound E may contain two double bonds. The ozonolysis reaction is



Since there is no loss of carbon atoms in an ozonolysis reaction, probably 2 molecules of formaldehyde are formed. In 2-ketopropanal, $-\underset{\text{O}}{\underset{\parallel}}{\text{C}}-$ and $-\text{CHO}$ groups must have resulted due to the cleavage of double bonds. Hence, the net




reaction can be explained as follows.



Hence, the compound E is 2-methyl-1, 3-butadiene.

Friedel Crafts Reaction

Friedel-Crafts Reaction



Identify the compound

An Organic compound A has 76.6% C and 6.38% H (Aspirin)

An organic compound A has 76.6% C and 6.38% H. Its vapour density is 47. It gives characteristic colour with FeCl_3 solution. The compound A when treated with CO_2 and NaOH at 140°C under pressure gives B which on being acidified gives C. The compound C reacts with acetyl chloride to give D which is a well known pain killer. Identify A, B, C and D and explain the reactions involved.

Solution The ratios of atoms present in one molecule of A are


$$\text{C} : \text{H} : \text{O} :: \frac{76.6}{12} : \frac{6.38}{1} : \frac{17.02}{16} :: 6.38 : 6.38 : 1.064 :: 6 : 6 : 1$$

Hence, Empirical formula of A is $\text{C}_6\text{H}_6\text{O}$. Molar empirical formula mass of A = 94 g mol^{-1}

This tallies with the given molar mass ($= 2 \times 47 \text{ g mol}^{-1}$). Thus Molecular formula of A is $\text{C}_6\text{H}_6\text{O}$.

Since, the compound A gives characteristic colour with FeCl_3 , it must be a phenol.

The given reactions are



Match the entries given on the left with those given on the right.

Friedel-Crafts	oil	alkenes
Fermentation	Lewis acid	soap
Dehydrohalogenation	cuprous chloride	anhydrous AlCl_3
Sandmeyer	yeast	chlorobenzene
Saponification	alcoholic alkali	ethanol

Ans :

Friedel-Crafts;	Lewis acid;	anhydrous AlCl_3
Fermentation;	yeast;	ethanol
Dehydrohalogenation;	alcoholic alkali;	alkenes
Sandmeyer;	cuprous chloride;	chlorobenzene
Saponification;	oil;	soap

Identify the compound

An organic compound A of molar mass 140.5 gm per mole has 68.32% Hydrogen

An organic compound A of molar mass 140.5 g mol^{-1} has 68.32% C, 6.4% H and 25.26% Cl. Hydrolysis of A with dilute acid gives compound B, $\text{C}_8\text{H}_{10}\text{O}$. Compound B can be oxidized under mild conditions to compound C, $\text{C}_8\text{H}_8\text{O}$. Compound C forms a phenylhydrazone D with PhNHNH_2 and gives a positive iodoform test. Deduce the structures of compounds A to D with proper reasoning.

Solution The ratios of atoms in a molecule of A are

$$\text{C} : \text{H} : \text{Cl} :: \frac{68.32}{12} : \frac{6.4}{1} : \frac{25.26}{35.5} :: 5.69 : 6.4 : 0.71 :: 8 : 9 : 1$$

Hence, Empirical formula of A is $\text{C}_8\text{H}_9\text{Cl}$. Molar empirical formula mass of A = 140.5 g mol^{-1}


This tallies with the given molar mass. Thus Molecular formula of A is $\text{C}_8\text{H}_9\text{Cl}$.

The high content of carbon indicates the presence of aromatic (or benzene) ring in the molecule.

Since Cl atom in the compound A is easily replaced during hydrolysis of A with dilute acid, the Cl atom must be attached to the aliphatic portion of the molecule.

Since the compound C gives a positive iodoform test, it must contain $-\text{COCH}_3$ group.

Since the compound C is obtained by mild oxidation of B, the latter must be a secondary alcohol. Hence, the compound A also contains secondary chlorine atom. Consistent with this, the given reactions are as follows.



Question

Bleaching powder and bleach solution are produced on a large scale and used in several household products. The effectiveness of bleach solution is often measured by iodometry.

25 mL of household bleach solution was mixed with 30 mL of 0.50 M KI and 10 mL of 4 N acetic acid. In the titration of the liberated iodine, 48 mL of 0.25 N $\text{Na}_2\text{S}_2\text{O}_3$ was used to reach the end point. The molarity of the household bleach solution is



- (a) 0.48 M (b) 0.96 M (c) 0.24 M (d) 0.024 M

Bleaching powder contains a salt of an oxoacid as one of its components. The anhydride of that oxoacid is

- (a) Cl_2O (b) Cl_2O_7 (c) ClO_2 (d) Cl_2O_6

Solutions :

For Q.11, we may write



$$\text{Amount of } \text{Na}_2\text{S}_2\text{O}_3 \text{ used} = VM = (48 \times 10^{-3} \text{ L}) (0.25 \text{ mol L}^{-1}) = 12 \times 10^{-3} \text{ mol}$$

$$\text{Amount of } \text{I}_2 \text{ generated} = \frac{1}{2} (12 \times 10^{-3} \text{ mol}) = 6 \times 10^{-3} \text{ mol}$$

Assuming 1 mol of household bleach produces 1 mol I_2 , we will have

$$\text{Amount of household bleach in 25 mL solution} = 6 \times 10^{-3} \text{ mol}$$

$$\text{Molarity of household bleach} = \frac{n}{V} = \frac{6 \times 10^{-3} \text{ mol}}{25 \times 10^{-3} \text{ L}} = 0.24 \text{ M}$$

For Q 12, we have

Bleaching powder contains $\text{Ca}(\text{OCl})\text{Cl}$.

The oxoacid of this salt is HOCl. The formation of this acid through its anhydride is $\text{Cl}_2\text{O} + \text{H}_2\text{O} \rightarrow 2\text{HOCl}$
anhydride oxoacid

Hence, we have

The **choice (c)** is correct

The **choice (a)** is correct.

Match the compounds/ions in Column I with their properties/reactions in Column II.

Column I

- (a) $\text{C}_6\text{H}_5\text{CHO}$
 (b) $\text{CH}_3\text{C} \equiv \text{CH}$
 (c) CN^-
 (d) I^-

Column II

- (p) gives precipitate with 2,4-dinitrophenylhydrazine
 (q) gives precipitate with AgNO_3
 (r) is a nucleophile
 (s) is involved in cyanohydrin formation

Ans :

$$(a) - (p), (q), (s); \quad (b) - (q); \quad (c) - (q), (r), (s); \quad (d) - (q)$$

In decreasing order of Oxidation state of Nitrogen

Which ordering of compounds is according to the decreasing order of the oxidation state of nitrogen ?

- (a) HNO_3 , NO, NH_4Cl , N_2 (b) HNO_3 , NO, N_2 , NH_4Cl
(c) HNO_3 , NH_4Cl , NO, N_2 (d) NO, HNO_3 , NH_4Cl , N_2

Solution :

The oxidation states of nitrogen in the given compounds are as follows.



Thus, the decreasing oxidation numbers of nitrogen is HNO_3 , NO, N_2 and NH_4Cl .

Therefore, the **choice (b)** is correct.

Gyan Question

The Kinetic energy of an electron in the second Bohr orbit

The kinetic energy of an electron in the second Bohr orbit of a hydrogen atom (a_0 is Bohr radius) is

- (a) $h^2/(4\pi^2ma_0^2)$ (b) $h^2/(16\pi^2ma_0^2)$ (c) $h^2/(32\pi^2ma_0^2)$ (d) $h^2/(64\pi^2ma_0^2)$

Solution :

Bohr model of an atom satisfies the following two requirements.

- Equality of centripetal and centrifugal forces, i.e.
- Quantization of angular momentum, i.e.

$$\frac{(Ze)(e)}{(4\pi\epsilon_0)r^2} = \frac{mv^2}{r} \quad mvr = n\left(\frac{h}{2\pi}\right)$$

Eliminating v in these two expressions, we get

$$r = n^2 \left[\frac{h^2}{4\pi^2 m (Ze^2 / 4\pi\epsilon_0)} \right] = \frac{n^2}{Z} a_0$$

where a_0 is the Bohr radius. The kinetic energy of the electron in Bohr orbit of an atom is

$$\text{KE} = \frac{1}{2}mv^2 = \frac{1}{2}m \left[\frac{n}{mr} \left(\frac{h}{2\pi} \right) \right]^2 = \frac{1}{2}m \left[\frac{n}{m(n^2 a_0 / Z)} \left(\frac{h}{2\pi} \right) \right]^2 = \frac{Z^2}{n^2} \left(\frac{h^2}{8\pi^2 m a_0^2} \right)$$


For the second Bohr orbit of hydrogen atom, $Z = 1$, and $n = 2$. Hence

$$\text{KE} = \frac{1}{32} \left(\frac{h^2}{\pi^2 m a_0^2} \right)$$

Therefore, the **choice (c)** is correct.

Question on Aldol reaction

The number of aldol reaction(s) that occurs in the given transformation



is

(a) 1




(b) 2

(c) 3

(d) 4

Solution :

The given reaction may be formulated as follows.



There are three aldol condensations and one Cannizzaro reaction.

Therefore, the **choice (c)** is correct

An organic compound undergoes first order decomposition

An organic compound undergoes first-order decomposition. The time taken for its decomposition to 1/8 and 1/10 of its initial concentration are $t_{1/8}$ and $t_{1/10}$, respectively. What is the value of $(t_{1/8}/t_{1/10}) \times 10$? ($\log 2 = 0.3$.)

Solution :

For first-order decomposition of A, the rate law is $\ln([A]/[A]_0) = -kt$

Hence $\ln(1/8) = -k t_{1/8}$ and $\ln(1/10) = -k t_{1/10}$

Thus $\frac{\ln(1/8)}{\ln(1/10)} = \frac{t_{1/8}}{t_{1/10}}$ or $\frac{\log 8}{\log 10} = \frac{t_{1/8}}{t_{1/10}}$

Hence $(t_{1/8}/t_{1/10}) = \log 8 = \log 2^3 = 3 \log 2 = 3 \times 0.3 = 0.9$ or $10(t_{1/8}/t_{1/10}) = 10 \times 0.9 = 9$


Therefore, the answer is **9**.

Allene has sp hybridization

In allene (C_3H_4), the type(s) of hybridization of the carbon atom is (are)

- (a) sp and sp^3 (b) sp and sp^2 (c) sp^3 only (d) sp^2 and sp^3


Solution :

The structure of allene is 

In this molecule, carbon atoms 1 and 3 are sp^2 hybridized while the carbon atom 2 is sp hybridized. Therefore, the **choice (b)** is correct.


Number of chiral carbon and optically active products

The number of optically active products obtained from the complete ozonolysis of the given compound



- (a) 0 (b) 1 (c) 2 (d) 4

Solution :



None of the products will be optically active. Therefore, the **choice (a)** is correct.





Carboxylic functional group is present in aspirin

The carboxylic functional group (—COOH) is present in

- (a) picric acid (b) barbituric acid (c) ascorbic acid (d) aspirin

Solution :


The structures of the given compounds are as follows.



Therefore, the **choice (d)** is correct.





Identify the compound

In the following reaction sequence, the compound J is an intermediate







$J(\text{C}_9\text{H}_8\text{O}_2)$ gives effervescence on treatment with NaHCO_3 and positive Baeyer's test.

Q - The compound K is




Q - The compound I is




Perkin Condensation

Solutions :

The compound J must contain a —COOH group as it gives effervescence with NaHCO_3 . Also, it is an unsaturated compound as it gives positive Baeyer's test (decolourization of pink colour of alkaline KMnO_4). Since the compound J is obtained by treating the compound I with $(\text{CH}_3\text{CO})_2\text{O}$ and CH_3COONa , and more over the compound J seems to be aromatic (high carbon content), the compound I amongst the choices given in Q.14 seems to be benzaldehyde as it shows Perkin condensation shown in the following




The conversion J to K is as follows.



Hence, we have
The **choice (c)** is correct.
The **choice (a)** is correct.

Question on isomers


Which of the given statement(s) about N, O, P and Q with respect to M is(are) correct?



- (a) M and N are non-mirror image stereoisomers. (b) M and O are identical
(c) M and P are enantiomers (d) M and Q are identical

Solution :

First of all, we may draw Fischer projections of the given compounds




From the Fischer projections, it follows that
M and N are non-mirror images.
M and P are enantiomers.

M and O are identical.
M and Q are not identical.

Therefore the **choices (a), (b) and (c)** are correct.


Aldohexose exists in D configuration

When the following aldohexose exists in D-configuration, the total number of stereoisomers in its pyranose form is



Solution:

In D-configuration, the hydroxyl group at the asymmetric carbon placed at the bottom of Fischer projection lies to the right of the straight skeleton. Thus, we will have




There are four chiral carbon atoms in the pyranose form of the given aldohexose. Thus, there will be $16 (= 2^4)$ stereoisomers, out of which 8 are of D-configurations (CH_2OH group above the ring) and 8 are of L-configurations (CH_2OH group below the ring).

Therefore, the correct answer is **8**.

Peptides

The substituents R_1 and R_2 for nine peptides are listed in the table given below. How many these peptides are positively charged at $\text{pH} = 7.0$?



Peptide	R_1	R_2
I	H	H
II	H	CH_3
III	CH_2COOH	H
IV	CH_2CONH_2	$(\text{CH}_2)_4\text{NH}_2$
V	CH_2CONH_2	CH_2CONH_2
VI	$(\text{CH}_2)_4\text{NH}_2$	$(\text{CH}_2)_4\text{NH}_2$
VII	CH_2COOH	CH_2CONH_2
VIII	CH_2OH	$(\text{CH}_2)_4\text{NH}_2$
IX	$(\text{CH}_2)_4\text{NH}_2$	CH_3

Solution

In general, the pH of the solution at which amino acids exist as zwitterion follows the order :

Acidic side chain < neutral chain < basic side chain

(pH = 3) (pH = 5.5 – 6.0) (pH = 9–10)

This pH is known as isoelectric point.

At $\text{pH} = 7$, the acidic side chain (for which isoelectric point < 7) exists as a negatively-charged species (e.g. $-\text{COO}^-$) and the basic side chain (for which isoelectric point > 7) exists as a positively charged species (e.g. $-\text{NH}_3^+$). Since peptides IV, VI, VIII and IX contain $-\text{NH}_2$ group in R_1 or/and R_2 group(s), these are expected to exist as positively-charged species.

Therefore, the correct answer is **4**.



Try to figure out what sequence is depicted

neopentane < isopentane < *n*-pentane. A branched chain isomer has a lower boiling point than a straight chain isomer. The more numerous the branches, the lower the boiling point. The branching lowers the boiling point for all families of organic compounds. Branching causes the shape of molecule to approach towards a sphere and thus surface area decreases, with the result that the intermolecular forces become weaker and are overcome at a lower temperature.


n-butyl alcohol < isobutyl alcohol < *sec*-butyl alcohol < *tert*-butyl alcohol. Polarity of O—H bond increases in the same order.



The presence of bulky groups near the site of reaction, whether in the alcohol or in the acid, slows down esterification as well as de-esterification (hydrolysis).



Try to figure out the sequences



adipic acid < succinic acid < malonic acid < oxalic acid

Acidity decreases with increase in the intervening CH_2 groups.

isobutane < *n*-butane < *n*-butyl chloride < *n*-butanol

chlorobenzene < benzene < toluene < methoxybenzene

Decreasing order of reactivity towards alcoholic silver nitrate:

2-bromo-1-phenylethene, α -phenylethyl bromide, β -phenylethyl bromide

Increasing order of reactivity towards aqueous NaOH:

chlorobenzene, *m*-chloronitrobenzene, *o*-chloronitrobenzene, 2,4-dinitrochlorobenzene,
2,4,6-trinitrochlorobenzene

Increasing reactivity towards HCN: CH_3CHO , CH_3COCH_3 , HCHO , $\text{C}_2\text{H}_5\text{COCH}_3$

Increasing basicity: *p*-toluidine, *N,N*-dimethyl-*p*-toluidine, *p*-nitroaniline, aniline.

Increasing ease of hydrolysis: $\text{CH}_3\text{COOC}_2\text{H}_5$, CH_3COCl , $(\text{CH}_3\text{CO})_2\text{O}$, CH_3CONH_2

Increasing order of acid strength:





ClCH_2COOH , $\text{CH}_3\text{CH}_2\text{COOH}$, $\text{ClCH}_2\text{CH}_2\text{COOH}$, $(\text{CH}_3)_2\text{CHCOOH}$, CH_3COOH

Increasing reactivity in nucleophilic substitution reactions: CH_3F , CH_3I , CH_3Br , CH_3Cl

major product H in the given reaction sequence


The major product H in the given reaction sequence

is $\text{CH}_3-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3 \xrightarrow[\text{HCN}]{\text{CN}^-} \text{G} \xrightarrow[\text{Heat}]{95\% \text{H}_2\text{SO}_4} \text{H}$




Solution :

The first reaction is nucleophilic addition reaction across the $-\text{C}=\text{O}$ bond.



In the second reaction, $-\text{CN}$ is hydrolysed to $-\text{COOH}$. The resultant molecule undergoes dehydration on heating.



Therefore, the **choice (a)** is correct.

Decreasing order of acidity of substituted benzoic acids:

p-chlorobenzoic acid, 2, 4-dichlorobenzoic acid, 2,4,6-trichlorobenzoic acid.

Decreasing order of acidity of carboxylic acids:

α -chlorophenylacetic acid, *p*-chlorophenylacetic acid, phenylacetic acid, α -phenylpropionic acid

Decreasing order of acidity of carboxylic acids:

p-nitrobenzoic acid, *p*-nitrophenylacetic acid, β -(*p*-nitrophenyl) propionic acid

Increasing order of basicity : ammonia, aniline, cyclohexylamine

Decreasing order of basicity: ethylamine, 2-aminoethanol, 3-amino-1-propanol

Decreasing order of basicity : aniline, *p*-methoxyaniline, *p*-nitroaniline

Increasing order of acidity: benzene sulphonic acid, benzoic acid, benzyl alcohol, phenol

Decreasing order of acidity: *m*-bromophenol, *m*-cresol, *m*-nitrophenol, phenol

Decreasing order of acidity of substituted phenol:

p-chlorophenol, 2, 4-dichlorophenol, 2, 4, 6-trichlorophenol

Decreasing order of reactivity towards KCN: benzyl chloride, chlorobenzene, ethyl chloride

Increasing order of nitration: benzene, chlorobenzene, nitrobenzene, toluene

Increasing order of reactivity towards alcoholic silver nitrate:

1-bromo-1-butene, 3-bromo-1-butene, 4-bromo-1-butene

Explain these by solving in right order -

Increasing reactivity towards S_N1 solvolysis:

benzyl chloride, *p*-chlorobenzyl chloride, *p*-methoxybenzyl chloride, *p*-methylbenzyl chloride,
p-nitrobenzyl chloride

Increasing order of reactivity towards elimination by alcoholic KOH:

1-phenyl-2-bromopropane, 1-phenyl-3-bromopropane

Decreasing order of reactivity towards aqueous HBr: Isomeric pentyl alcohols

Increasing order of reactivity towards aqueous HBr:

1-phenyl-1-propanol, 3-phenyl-1-propanol, 1-phenyl-2-propanol

Decreasing order of reactivity towards aqueous HBr:

benzyl alcohol, *p*-cyanobenzyl alcohol, *p*-hydroxybenzyl alcohol

Increasing order of reactivity towards aqueous HBr:

benzyl alcohol, diphenylmethanol, methanol and triphenylmethanol

Decreasing order of reactivity towards bromination:

anisole, benzene, toluene, chlorobenzene, nitrobenzene, phenol

Increasing order of reactivity towards bromination:

hydroquinone, *p*-methoxyphenol, *p*-methylphenol, *p*-chlorophenol, *p*-nitrophenol, *sym*-trihydroxybenzene

Decreasing order of acidity of carboxylic acids:

butanoic acid, 2-bromobutanoic acid, 3-bromobutanoic acid, 4-bromobutanoic acid

Decreasing reactivity towards S_N1 substitution: 1-chloropropene 3-chloropropene, *n*-propylchloride

Decreasing acidity: H_2O , $HC\equiv CH$, NH_3 , RH , ROH

Decreasing basicity: R^- , $HC\equiv C^-$, NH_2^- , OH^- , OR^-

Decreasing order of reactivity towards the addition of HCl:

styrene, *p*-chlorostyrene, *p*-methylstyrene, *p*-nitrostyrene

Decreasing order of reactivity towards dehydration:




α -phenyl ethyl alcohol, α -(*p*-nitrophenyl) ethyl alcohol, α -(*p*-aminophenyl) ethyl alcohol

Various Increasing Properties

Increasing stability of free radicals: $\bullet\text{CH}_3$, 1° , 2° , 3° , allyl, vinyl

Increasing stability of carbocation: CH_3^+ , 1° , 2° , 3°

Increasing enthalpy of reaction:




Increasing order of reactivity towards $\text{S}_{\text{N}}2$ displacement:

1-bromopentane, 2-bromopentane, 2-bromo-2-methylbutane


compound undergoes decarboxylation most readily

The compound that undergoes decarboxylation most readily under mild condition is



Solution :

β -Ketocarboxylic acid is unstable acid. It readily undergoes decarboxylation through a cyclic transition state.



Therefore, the **choice (b)** is correct.

A compound contains 88.89% C and 11.1% H. It gives white precipitate with ammoniacal silver nitrate. The compound is


- (a) $\text{C}_3\text{H}_7\text{C}\equiv\text{CH}$ (b) $\text{C}_8\text{H}_9\text{C}\equiv\text{CH}$ (c) $\text{C}_6\text{H}_{11}\text{C}\equiv\text{CH}$ (d) $\text{C}_4\text{H}_7\text{C}\equiv\text{CH}$

Ans :

Empirical formula contains $C : H :: \frac{88.89}{12} : \frac{11.1}{1} :: 7.4 : 11.11 :: 2 : 3$


Only the compound $C_6H_{11}C \equiv CH$ has composition 4 times the composition of empirical formula.

More the number of electron withdrawing group stronger is the acid








The more the number of electron-withdrawing group, the stronger the acid.

Try to figure out what sequence is depicted



The more the electronegativity of halogen, the stronger the acid.



$(CH_3)_3N < CH_3NH_2 < (CH_3)_2NH$. $(CH_3)_3N$ is the weakest because of steric factors.

1-butanol < 2-butanol < 2-methyl-2-propanol. A tertiary alcohol reacts immediately, a secondary alcohol reacts within five minutes and a primary alcohol does not.

2-methyl-2-propanol < 2-butanol < 1-butanol. Reactivity of ROH is $CH_3OH > 1^\circ > 2^\circ > 3^\circ$.

An organic compound A contains 49.32% C, 9.59% H and 19.18% N and rest oxygen. The compound A on boiling with NaOH gives nitrogen free carboxylic acid whose silver salt contains 59.67% Ag. The compound A is

- (a) CH_3CONH_2 (b) $C_2H_5CONH_2$ (c) $C_3H_7CONH_2$ (d) $C_3H_5CONH_2$

Ans :


An organic compound A contains 49.32% C, 9.59% H and 19.18% N and rest oxygen. The compound A on boiling with NaOH gives nitrogen free carboxylic acid whose silver salt contains 59.67% Ag. The compound A is

- (a) CH_3CONH_2 (b) $C_2H_5CONH_2$ (c) $C_3H_7CONH_2$ (d) $C_3H_5CONH_2$

Hence $\frac{108}{M_R + 44 + 108} = 0.5967$. This gives $M = 29$ i.e. $R \equiv C_2H_5$.

Hence, the compound is $C_2H_5CONH_2$.


Try to figure out what sequence is depicted



The smaller the alkyl group, the more reactive the carbonyl group.




The more stable the intermediate carbocation, the greater the reactivity.



Conjugated dienes form the more stable allyl carbocations and are thus more reactive than alkenes. Alkyl groups on the unsaturated carbon atoms increase reactivity.



tert-butanol < *sec*-butanol < *n*-butanol < CH₃OH. Alkyl group makes an alcohol less acidic. Bigger the alkyl group, the less acidic the alcohol. Methanol is the strongest and tertiary alcohols are the weakest.




An aromatic compound contains 69.4% C and 5.8% H

An aromatic compound contains 69.4% C and 5.8% H. A sample of 0.3 g of this compound give ammonia which neutralizes 25 mL of 0.05 M H₂SO₄. The empirical formula of the compound is

- (a) C₅H₅N₂O (b) C₇H₇NO (c) C₆H₅NO₂ (d) C₆H₈NO

Ans :

$$\text{Amount of H}_2\text{SO}_4 \text{ neutralized} = (25 \times 10^{-3} \text{ L})(0.05 \text{ M}) = 1.25 \times 10^{-3} \text{ mol}$$



$$\text{Amount of NH}_3 \text{ evolved} = 2 \times 1.25 \times 10^{-3} \text{ mol} = 2.5 \times 10^{-3} \text{ mol}$$

$$\text{Mass of N in the compound} = (2.5 \times 10^{-3} \text{ mol}) (14 \text{ g mol}^{-1}) = 0.035 \text{ g}$$

$$\text{Per cent of N in the compound} = \frac{0.035}{0.3} \times 100 = 11.67$$

$$\text{Per cent of O in the compound} = 100 - (69.4 + 5.8 + 11.67) = 13.13$$


Ratio of atoms in the compound is

$$\text{C} : \text{H} : \text{N} : \text{O} :: \frac{69.4}{12} : \frac{5.8}{1} : \frac{11.67}{14} : \frac{13.13}{16} :: 5.78 : 5.8 : 0.83 : 0.82 :: 7 : 7 : 1 : 1$$


Hence, empirical formula: C₇H₇NO

Alternatively, calculate per cent of C in the given choices which comes out to be (a) 0.55, (b) 0.69, (c) 0.58 and (d) 0.65 only for choice (b), the answer tallies.

Zwitter Ions - The Hydrogen atom (rather ion) moves to different positions depending on the PH (So can have a positive charge at some position, or a negative charge at some other position, depending on the pH)




Electrophoresis - is a method to confirm if a positive charge is present or a negative charge is present in the ion.




Iso -Electric point - the pH at which the ion (molecule) does not migrate to any electrodes.

What happens if oxalic acid is heated with conc sulphuric acid ?



We get Carbon monoxide




IMPORTANT ORDER AND FACTS OF ORGANIC CHEMISTRY





1. $\text{RCOCl} > \text{RCOOCOR} > \text{RCOOR} > \text{RCONH}_2$ Nucleophilic substitution reaction.
2. $\text{HI} > \text{HBr} > \text{HCl} > \text{RCOOH} > \text{C}_6\text{H}_5\text{OH} > \text{H}_2\text{O} > \text{CH} \equiv \text{CH} > \text{NH}_3$ (Acidic nature).

3. $\text{CCl}_3\text{CHO} > \text{HCHO} > \text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_3$ Nucleophilic addition reaction.


4. $\text{CH}_2 = \text{CH}_2 > \text{CH} \equiv \text{CH} > \text{C}_6\text{H}_6$ Electrophilic addition reaction.



Substitution Reaction



10. $(\text{CH}_3)_2\text{C} = \text{C}(\text{CH}_3)_2 > \text{CH}_3 - \text{CH} = \text{C} - (\text{CH}_3)_2 > \text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_3 > \text{CH}_3 - \text{CH} = \text{CH}_2$ (Stability)



12. $\text{NH}_2^{\ominus} > \text{OH}^{\ominus} > \text{NH}_3 > \text{H}_2\text{O}$ (Basic strength)

13. $\text{NI}_3 > \text{NBr}_3 > \text{NCl}_3 > \text{NF}_3$ (Basic strength)

14. $\text{Br}_2 > \text{Cl}_2 > \text{I}_2$ (Selectivity for halogenation)

15. Halogenation of alkenes by cyclic halonium state, so anti attack takes place.

16. Hydroboration followed by oxidation is always anti markownikoff's addition due to steric effect.

17. Oximercuration - demercuration is markownikoff's addition of water because some carbocation character in cyclic mercurium state.

18. CHCl_3 in the presence of strong bases forms biradical $:\text{CCl}_2$ which undergo addition with double or triple bonds.

19. When conjugated diene reacts with alkene or alkyne it is known as diel's elder synthesis.

20. Ozonolysis of cyclo alkene forms one mole dialdehyde while ozonolysis of cyclo alkadiene forms two moles of dialdehyde.

21. Ozonolysis with $(\text{CH}_3)_2\text{S}$ is known as reductive ozonolysis.

22. Hydration of alkyne occur's in HgSO_4 and dil H_2SO_4 .

23. 1-alkynes forms ppt with $\text{Ag}(\text{NH}_3)_2^\ominus$ and $\text{Cu}(\text{NH}_3)_4^+$.

24. Cis-2-butene reacts with Br_2 to forms dl(±) pair of enantiomers of 2,3-dibromobutane while in case of trans-2-butene forms meso-2,3-dibromo butane due to anti addition always.

25. Haloform test given by species with CH_3CO -group but not in case of A.A.E. and tert. Butyl alcohol.

26. Chloral reacts with chloro benzene in con. H_2SO_4 to form insecticide DDT.

27. NBS is used for free radical allylation.

28. Rate for S_N^1 reaction is $3^\circ > 2^\circ > 1^\circ$ in protic polar solvent.

29. Rate for S_N^2 reaction is $1^\circ > 2^\circ > 3^\circ$ in polar aprotic solvent like DMSO, DMF, HMPT.

30. Chemical reactions like Hoffmann carbylamine and Reimer Tiemann's reaction active species is biradical CCl_2 .


31. If cyclo 1,3-penta diene reacts with CHCl_3 and potassium tert. butoxide to form chlorobenzene.

32. Alkyl halides reacts with AgCN to form isocyanides due to ambident nature of nucleophile, other ambident nucleophiles are NO_2^\ominus and SO_3^{-2} .

33. In dehydration of alcohols active species is carbocation so rearrangement occurs like hydride shift or alkyl shift.

34. Dehydration of cyclobutyl methyl alcohol ring expansion takes place, formation of cyclo pentene occurs.

35. In esterification where acid reacts with alcohol to form ester, - OH given by acid while - H by alcohol



36. Ether's reacts with HI to form alcohol and halide where fission of lower ether by S_N^2 mechanism while higher ethers like ter. butyl methyl ether or alkyl methyl ether by S_N^1 mechanism.

37. Quantitative estimation of ethers is done by ziesel's method.

38. If unsym. cyclic ether undergo fission it depends upon medium whether it is acid or basic like in acidic medium some character of carbocation so nucleophile goes to carbon where more alkyl groups are there while reverse in basic medium due to steric factor.

39. Aldehydes are reducing agent while ketones are not.

40. Aldehydes and ketones are separated by tollen's reagent.

41. Carbonyl and noncarbonyl are separated by sodium bisulphite and bradye's reagent.

42. As the size of alkyl group increases steric hindrance comes into play, reactivity towards nucleophilic addition decreases.

43. Aldehydes with α -hydrogen atom in the presence of dil base undergo enolization and form α




carbonian to give aldol product.











44. If there is two-CHO group with α -hydrogen atom to form cyclic intra aldol product.
45. Aldehydes without α -hydrogen atom in the presence of con. alkali to form each molecule of acid and alcohol by hydride active species.
46. Glyoxal reacts with con. KOH to form glycolate ion by Intra Cannizaro's.
47. Pinacol pinacolone type reactions involve protonation, deprotonation and alkyl shift.
48. Aldol, Perkin, reformatsky and knovengel's reactions are classified as carbanian active reaction.
49. In Beckmann's rearrangement migration of group which is anti to-OH group takes place.
50. Beckmann's rearrangement is a reaction of oximes in the presence of H_2SO_4 or PCl_5 to produce N-alkyl amide derivatives.
51. Migratory attitude of alkyl group in Pinacol-pinacolone, beckmann's and bayer villegar oxidation is $-C_6H_5 > (CH_3)_3C - > (CH_3)_2CH - > C_2H_5 - > CH_3-$
52. Cyclo hexanoneoxime on beckmann's reaction gives caprolectum which on reaction with H^+ to give polymer nylon-6
53. 2-methyl propanal even contains α -hydrogen atom but does not give aldol reaction.
54. Diphenyl glyoxal reacts with con. KOH to form salt of benzillic acid.
55. Propanone in the presence of dry HCl gas by enolic intermediate to form diacetone alcohol undergo heating form mesityl oxide. If this again reacts with propanone to form phoron.
56. Benzaldehyde reacts with alc. KCN to form Benzoin which on oxidation form benzil.
57. By wittig reaction carbonyl compounds are converted into E-Z form of alkene.
58. Benzaldehyde do not respond to benedict's and fehling's solution due to less reducing power of aromatic aldehyde.
59. HCOOH respond to oxidising agent due to presence of -CHO group.
60. $HI > H_2SO_4 > HNO_3 > RCOOH > H_2CO_3 > C_6H_5OH > H_2O > CH \equiv CH > NH_3 > CH_3 - C \equiv CH > CH_2 = CH_2 > H_2$ this is decreasing acidic nature.
61. Acids with α -hydrogen atom when reacts with halogen in the presence of P to form α -haloacid (HVZ).
62. Carboxylic acid on reaction with PCl_5 , CH_2N_2 and H_2O to form higher acid or next homologue.
63. For reactivity of acid derivatives use funda weaker the base better the leaving group.
64. Acid amide on reaction with PCl_5 to form alkane nitrile.
65. Anhydride on reaction with carbonyl compound in the presence of base (carbanian) forms α, β -unsaturated carboxylic acid (perkin reaction)







66. Amide on reaction with Br_2 and alkali to form primary amine of lower homologue. Intermediate species is nitrene which undergo intra rearrangement to form RNH_2 .
67. Ester's with α -hydrogen atom in the presence of strong base to form carbanion undergo nucleophilic substitution reaction forms β -keto ester for example ethyl acetate in the presence of pot. ter. Butoxide form aceto acetic ester (AAE). reaction is known as clesen's ester condensation of four types
- Simple clesen's ester condensation.
 - CROSS clesen's ester condensation.
 - Intra CEC (Dieckmann's condensation).
 - mixed clesen ester condensation.
68. For a compound to be aromatic it must be cyclic, planar and obey $(4n + 2) \pi$ electron rule.
69. Cyclooctatetraene is non aromatic compound while pyrrole, pyredene, furan, cyclopentadieneylanion all are aromatic.
70. In aromatic electrophilic substitution reaction there is no hydrogen isotopic effect except sulphonation and iodination.
71. m- directing groups like nitrobenzene and benzaldehyde cannot undergo fridal craft reaction.
72. Phenol is less acidic than general carboxylic acid, cannot react with NaHCO_3 .
73. Anilene is more reactive than phenol towards electrophilic substitution reaction because less energy difference between nitrogen and carbon.
74. Benzene diazonium chloride reacts with phenol or anilene to form azo compounds.
75. $\text{C}_6\text{H}_5\text{NH}_2 > \text{C}_6\text{H}_5\text{OH} > \text{C}_6\text{H}_5\text{OR} > \text{C}_6\text{H}_5\text{Br} > \text{C}_6\text{H}_5\text{NO}_2$ (Reactivity towards electrophilic substitution reaction)


General Series - Chemistry Fact Sheet - 3

General series	Order	Why ?
1. B.P. of $\text{CH}_3\text{CH}_2\text{OH}$, CH_3OCH_3 , CH_3CH_3 I II III	I > II > III	There is intermolecular H-bonding I. III has weak force of attraction and is most volatile. Intramolecular H-bonding in o-isomer makes it more volatile.
2. B.P. of o, m, p-nitro phenol	o < m < p	

3. **Reactivity of ... with Tollen's reagent**
 HCHO , CH_3CHO , CH_3COCH_3 , $\text{C}_6\text{H}_5\text{CHO}$
 I II III IV
 I > II > IV > III
 -CHO group is easily oxidised compared to keto group due to reducing hydrogen.
4. **Reactivity of ... with Fehling's solution**
 I II III IV (above)
 I > II > IV > III
 -do-
5. **Extent of hydration of**

 (I) (II) (III) (IV)
 I < II < III < IV
 Aldehydes are more hydrated than ketones. Halide makes C of carbonyl group more electropositive.
6. **Electrophilic nature of for nucleophilic attack**

 I II III
 I > II > III
 CH₃ group decreases +ve charge on C hence nucleophilic attack.
7. **Reactivity of isomeric 1°, 2°, 3° butyl halide towards elimination (E1 or E2)**
 $3^\circ < 2^\circ < 1^\circ$
 due to stability of intermediate carbocation
8. **Dehydration of**

 (I) (II) (III) (IV)
 IV < I < II < III
 Alcohol leading to increase in conjugation due to dehydration is more easily dehydrated. IV is vinylic, hence least.
9. **Stability of**

- $\text{CH}_2=\text{CH}_2$, $\text{CH}_3\text{CH}=\text{CH}_2$, $\text{CH}_3\text{CH}=\text{CHCH}_3$
 I II III
 $\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{CHCH}_2 \\ \diagup \\ \text{CH}_3 \end{array}$ $\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{CHCH}_3 \\ \diagup \\ \text{CH}_3 \end{array}$
 IV V
 $\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{C} \\ \diagup \\ \text{CH}_3 \end{array}$ $\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{C} \\ \diagup \\ \text{CH}_3 \end{array}$
 VI
- Substituted alkenes are more stable. More the alkyl groups are attached to the doubly bonded carbon atom more is the stability.
- I < II < III < IV < V < VI
10. **Stability of**
- (I)  (II)  (III) 
- I < III < II
11. **Stability of**
- (I)  (II)  (III)  (IV) 
- III > II > I > IV
12. **Stability of**
- I $\text{CH}_2=\overset{\bullet}{\text{C}}\text{H}$, II $\overset{\bullet}{\text{C}}\text{H}_2\text{CH}=\text{CH}_2$, (III)  (IV) 
- I < IV < II < III is 3° allylic and II is 1° allylic
13. **Dehydration of**
1°, 2°, 3° isomeric butyl alcohol
- 3° < 2° < 1°
14. **Boiling points of**
n-butyl amine, *n*-butyl alcohol, *n*-pentane
I II III
- II > I > III
15. **Formation of**
- I  II $\text{R}_3\overset{\bullet}{\text{C}}$, III $\text{R}_2\overset{\bullet}{\text{C}}\text{H}$, IV $\text{R}\overset{\bullet}{\text{C}}\text{H}_2$
- I > II > III > IV
16. **Reactivity of C–H bond (abstraction of H)**
- greater the stability of intermediate, greater the reactivity of chemical reaction. I, II have H-bonding but electronegativity of O > N hence H-bonding in II > I greater the stability, easier the formation of particular species.






atoms separating the ring from the positively charged substituent increases, deactivating effect decreases due to less electronegativity.

$-\text{O}^\ominus$ is best able to donate electrons there by giving a very stable uncharged intermediate. In

22.



II > I > III

$\text{CH}_3\text{CO}-$ cross conjugation diminished its ability to donate electrons to an arenium ion. Intermediates are benzylic cations. So CH_3O (electron repelling) gives greater stability through delocalisation while NO_2 (electron attracting) decreases stability.

23.

Relative reactivity of ... towards $\text{S}_\text{N}1$ reaction
benzyl chloride, *p*-methoxy benzyl chloride
and *p*-nitro benzyl chloride

I II III

II > I > III

delocalisation while NO_2 (electron attracting) decreases stability.

24.

Relative reactivity of ... towards $\text{S}_\text{N}1$ and $\text{S}_\text{N}2$ reaction
 PhCH_2Cl PhCHClMe PhCClMe_2
(I) (II) (III)
(Ph stands for phenyl, C_6H_5)

$\text{S}_\text{N}1$:
III > II > I
 $\text{S}_\text{N}2$:
II < II < I


$\text{S}_\text{N}1$: $1^\circ < 2^\circ < 3^\circ$
alkyl halide
 $\text{S}_\text{N}2$: $3^\circ < 2^\circ < 1^\circ$
alkyl halide

25.

Relative reactivity of ... with E^+ (electrophile) in S_E reaction.



II > I > III

$-\text{NO}_2$ deactivates benzene ring for S_E



26.

Order of S_N2 reactivity of alkoxide nucleophiles









$I < IV < V < III < II$

S_N2 reactivity is susceptible to steric hindrance by the nucleophile as well as by the size of alkyl group.




Basic power - Chemistry Fact Sheet - 2



	Basic power	Order	Why ?
1.		I > III > II	lone pair on N is not used in resonance of π -electrons in I. In II lone pair of the ring is itself used in delocalisation while that of outside ring in III.
2.		I > II > III > IV	$-\text{OCH}_3$ is strong electron donating group. This is due to ortho effect , all the aniline are less basic than <i>p</i> -substituted aniline due to steric hindrance .
3.		I > II > III > IV	I (hyper conjugation and induction) II (induction) IV (ortho effect), ortho effect normally decreases basic nature.

4.  II > I > III
- In II there is sp^3 hybridised C, In I, sp^2 . NO_2 is electron withdrawing.
5.  III > I > II
- lone pair on N is used in delocalisation of π -electrons in aromatic amines while cyclohexyl is electron repelling (III); in II, lone pair on N is used by two benzene ring. NO_2 is electron-withdrawing, thus nitro-anilines are less basic than aniline. IV is less basic than III because $-\text{NO}_2$ is closer and exerts a stronger inductive effect.
6.  I > II > III > IV
7.  III > I > II
- phenyl and $-\text{COCH}_3$ are electronwithdrawing and $-\text{C}_6\text{H}_5 < \text{COCH}_3$
8.  I < II < III
- Electron donating nature of $\text{C}_2\text{H}_5 > \text{CH}_3$ So more basic strength.



Acidic Powers & Their Orders - Chemistry Fact Sheet

	Acidic power	Order	Why ?
1.	1-, 2-, 3- chlorobutanoic acid I II III		Farther the (-I) group (Cl), lesser the acidic strength III < II < I
2.	1-, 2-, 3- methyl pentanoic acid I II III		Farther the (+I) group, greater the acidic power I < II < III
3.			-CH ₃ is electron donating and -NO ₂ is electron attracting II < I < III
4.			-CH ₃ is electron repelling; decreases acidic strength of phenol II < I < III
5.			-OCH ₃ group contains +M effect and decreases acidic power. III < I < II

6.  (I) (II) (III) $I < II < III$
7.  (I) (II) (III) $I > II > III$
8. oxalic acid, succinic acid, malonic acid, adipic acid
I II III
adipic acid (all dibasic) IV $I > II > III > IV$
9. *o*- *m*- *p*- nitrobenzoic acid
I II III $II < III < I$
10. *o*- *m*- *p*- hydroxy benzoic acid
I II III $III < II < I$
- NO₂ is electron attracting; III is more resonance stabilised than I and also than II. In I, only inductive effect is operative. sp² hybridised carbon of I, II are more electronegative hence acid strength is increased. Benzylic (C₆H₅CH₂) is more stabilised than allylic (CH₂=CHCH₂). Effect of one –COOH on the other decreases as its distance between them increases, (COOH)₂ is maximum acidic. –NO₂ is electron attracting (-I effect) –OH shows electron withdrawing nature at *o*- and *m*- and electron repelling at *p*-, *o*- isomer due to intramolecular bonding in salicylate ion is stronger than *m*- isomer

Silica is soluble in which of the following acids ?

(1) HF (2) HCl (3) HBr (4) HI

Solution : $\text{SiO}_2 + 4\text{HF} \rightarrow \text{SiF}_4 + 2\text{H}_2\text{O}$ and again $\text{SiF}_4 + 2\text{HF} \rightarrow \text{H}_2\text{SiF}_6$ (Hydrofluorosilicic acid)

$\text{K}_3\text{Co}(\text{NO}_2)_6$ is Fischer's Salt



The brown ring test for NO_3^- is due to formation of the complex $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]^{2+}$

Merck's perhydrol is 30.4% H_2O_2

Elements having one or more solvated ions in water

1 IA	Elements having one or more solvated ions in water																VIII 18
H	IIA											IIIA 13	IVA 14	VA 15	VIA 16	VIIA 17	He
Li	Be											B	C	N	O	F	Ne
Na	Mg	IIIB 3	IVB 4	VB 5	VIB 6	VIB 7	VIII 8	VIII 9	VIII 10	IB 11	IIB 12	Al	Si	P	S	Cl	Ar
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Cs	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
Rb	Ba	lan	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
Fr	Ra	act	Rf	Db	Sg	Bh	Hs	Mt	Ds								
												cations anions					

H_2SO_5 , $\text{H}_2\text{S}_2\text{O}_8$ have peroxide linkages





1 H																	2 He	
		 Ferromagnetic Antiferromagnetic																
		 Paramagnetic Diamagnetic																
3 Li	4 Be											5 B	6 C	7 N	8 O	9 F	10 Ne	
11 Na	12 Mg											13 Al	14 Si	15 P	16 S	17 Cl	18 Ar	
19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr	
37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe	
55 Cs	56 Ba	57 La	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn	
87 Fr	88 Ra	89 Ac																
			58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu		

Elements & Country of Discovery



1 H																	2 He	
3 Li	4 Be											5 B	6 C	7 N	8 O	9 F	10 Ne	
11 Na	12 Mg											13 Al	14 Si	15 P	16 S	17 Cl	18 Ar	
19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr	
37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe	
55 Cs	56 Ba	57 La	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn	
87 Fr	88 Ra	89 Ac																
			58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu		

Bromine reacting with NaOH in cold and hot give different mix of products. Specify






Malachite decomposes to give A + CO₂ + H₂O and compound A on reduction with Carbon gives CO + B

Identify A and B





-

Some examples of Complex Anions



1 H																	2 He																														
3 Li	4 Be											5 B	6 C	7 N	8 O	9 F	10 Ne																														
11 Na	12 Mg											13 Al	14 Si	15 P	16 S	17 Cl	18 Ar																														
19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr																														
37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe																														
55 Cs	56 Ba	57 La	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn																														
87 Fr	88 Ra	-103	104 Rf	105 Db	106 Sg	107 Bh	108 Hs	109 Mt	110 Ds	111 Rg	112 Cn	113 Uut	114 Fl	115 Uup	116 Lv	117 Uus	118 Uuo																														
<table border="1"> <tbody> <tr> <td>57 La</td> <td>58 Ce</td> <td>59 Pr</td> <td>60 Nd</td> <td>61 Pm</td> <td>62 Sm</td> <td>63 Eu</td> <td>64 Gd</td> <td>65 Tb</td> <td>66 Dy</td> <td>67 Ho</td> <td>68 Er</td> <td>69 Tm</td> <td>70 Yb</td> <td>71 Lu</td> </tr> <tr> <td>89 Ac</td> <td>90 Th</td> <td>91 Pa</td> <td>92 U</td> <td>93 Np</td> <td>94 Pu</td> <td>95 Am</td> <td>96 Cm</td> <td>97 Bk</td> <td>98 Cf</td> <td>99 Es</td> <td>100 Fm</td> <td>101 Md</td> <td>102 No</td> <td>103 Lr</td> </tr> </tbody> </table>																		57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu	89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr
57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu																																	
89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr																																	
<p>Known in antiquity</p> <p>also known when (akw) Levoisier published his list of elements (1789)</p> <p>akw Mendeleev published his periodic table (1869)</p> <p>akw Deming published his periodic table (1923)</p>						<p>akw Seaborg published his periodic table (1945)</p> <p>also known (ak) up to 2000</p> <p>ak to 2012</p>																																									



The ionization potential of Pb is higher than Sn due to poor shielding by 14f electrons present in Pb

Selenous acid H_2SeO_3 (Oxidation Number of Se is +4)

Telluric acid H_6TeO_6 (Oxidation Number of Te is +6)

Electronegativity Values (in Pauling Scale)

H																	He
2.1																	—
Lr	Be											B	C	N	O	F	Ne
1.0	1.5											2.0	2.5	3.0	3.5	4.0	—
Na	Mg											Al	Si	P	S	Cl	Ar
0.9	1.2											1.5	1.8	2.1	2.5	3.0	—
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
0.8	1.0	1.3	1.5	1.6	1.6	1.5	1.8	1.8	1.8	1.9	1.6	1.6	1.8	2.0	2.4	2.8	—
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
0.8	1.0	1.2	1.4	1.6	1.8	1.9	2.2	2.2	2.2	1.9	1.7	1.7	1.8	1.9	2.1	2.5	—
Cs	Ba	La-Lu	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
0.7	0.9	1.1-1.2	1.3	1.5	1.7	1.9	2.2	2.2	2.2	2.4	1.9	1.8	1.8	1.9	2.0	2.2	—

Microcosmic Salt is $Na(NH_4)HPO_4$ Sodium ammonium hydrogen phosphate


Which of the following electrolyte will be most effective in coagulation of gold sol ?

- (1) NaNO_3 (2) $\text{K}_4\text{Fe}(\text{CN})_6$ (3) Na_3PO_4 (4) MgCl_2


Gold sol is a negatively charged coagulation which is carried out by positively charged ion furnished by electrolyte

Coagulating power depends on the valency of cation of electrolyte. (Schulze Hardy Law) Mg^{2+} ion has highest valency so MgCl_2 is most effective.

Organometallic compounds are Metal atoms directly linked with Carbon. So $\text{Ti}(\text{OC}_3\text{H}_7)_4$ is not an organometallic compound.



V.T.E																		
Periodic table of electronegativity using the Pauling scale																		
→ Atomic radius decreases → Ionization energy increases → Electronegativity increases →																		
Group →	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
↓ Period																		
1	H 2.20																	He
2	Li 0.98	Be 1.57											B 2.04	C 2.55	N 3.04	O 3.44	F 3.98	Ne
3	Na 0.93	Mg 1.31											Al 1.61	Si 1.90	P 2.19	S 2.58	Cl 3.16	Ar
4	K 0.82	Ca 1.00	Sc 1.36	Ti 1.54	V 1.63	Cr 1.66	Mn 1.55	Fe 1.83	Co 1.88	Ni 1.91	Cu 1.90	Zn 1.65	Ga 1.81	Ge 2.01	As 2.18	Se 2.55	Br 2.96	Kr 3.00
5	Rb 0.82	Sr 0.95	Y 1.22	Zr 1.33	Nb 1.6	Mo 2.16	Tc 1.9	Ru 2.2	Rh 2.28	Pd 2.20	Ag 1.93	Cd 1.69	In 1.78	Sn 1.96	Sb 2.05	Te 2.1	I 2.66	Xe 2.60
6	Cs 0.79	Ba 0.89	*	Hf 1.3	Ta 1.5	W 2.36	Re 1.9	Os 2.2	Ir 2.20	Pt 2.28	Au 2.54	Hg 2.00	Tl 1.62	Pb 1.87	Bi 2.02	Po 2.0	At 2.2	Rn 2.2
7	Fr 0.7	Ra 0.9	**	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Uut	Fl	Uup	Lv	Uus	Uuo
* Lanthanoids	La 1.1	Ce 1.12	Pr 1.13	Nd 1.14	Pm 1.13	Sm 1.17	Eu 1.2	Gd 1.2	Tb 1.1	Dy 1.22	Ho 1.23	Er 1.24	Tm 1.25	Yb 1.1	Lu 1.27			
** Actinoids	Ac 1.1	Th 1.3	Pa 1.5	U 1.38	Np 1.36	Pu 1.28	Am 1.13	Cm 1.28	Bk 1.3	Cf 1.3	Es 1.3	Fm 1.3	Md 1.3	No 1.3	Lr 1.3			




Periodic Table of the Elements
Abundance of Elements
in Seawater near the Surface

all values are in mg/L.

1A	2A	3B	4B	5B	6B	7B	8B	1B	2B	3A	4A	5A	6A	7A	8A		
1 H 10000 Hydrogen	4 He 1×10^6 Helium	3 B 4.4 Boron	5 Li 1.5×10^6 Lithium	6 C 78 Carbon	7 N 1.5×10^6 Nitrogen	8 O 85000 Oxygen	9 F 7.5 Fluorine	10 Ne 1.5×10^6 Neon	11 Na 10800 Sodium	12 Mg 1200 Magnesium	13 Al 2×10^6 Aluminum	14 Si 72 Silicon	15 P 1.5×10^6 Phosphorus	16 S 900 Sulfur	17 Cl 19400 Chlorine	18 Ar 1.5×10^6 Argon	
19 K 190 Potassium	20 Ca 411 Calcium	21 Sc 1.5×10^6 Scandium	22 Ti 1×10^6 Titanium	23 V 2.5×10^6 Vanadium	24 Cr 2×10^6 Chromium	25 Mn 2×10^6 Manganese	26 Fe 2×10^6 Iron	27 Co 2×10^6 Cobalt	28 Ni 1.5×10^6 Nickel	29 Cu 2.5×10^6 Copper	30 Zn 4×10^6 Zinc	31 Ga 2×10^6 Gallium	32 Ge 1.5×10^6 Germanium	33 As 1.5×10^6 Arsenic	34 Se 2×10^6 Selenium	35 Br 11.2 Bromine	
37 Rb 1.5×10^6 Rubidium	38 Sr 7.5 Strontium	39 Y 1×10^6 Yttrium	40 Zr 2×10^6 Zirconium	41 Nb 1×10^6 Niobium	42 Mo 1×10^6 Molybdenum	43 Tc Technetium	44 Ru 1×10^6 Ruthenium	45 Rh Rhodium	46 Pd Palladium	47 Ag 1×10^6 Silver	48 Cd 1×10^6 Cadmium	49 In 1×10^6 Indium	50 Sn 4×10^6 Tin	51 Sb 2×10^6 Antimony	52 Te 1.5×10^6 Tellurium	53 I 10000 Iodine	
55 Cs 1×10^6 Cesium	56 Ba 1.5×10^6 Barium	57-71 Lanthanides	72 Hf 2×10^6 Hafnium	73 Ta 2×10^6 Tantalum	74 W 1×10^6 Tungsten	75 Re 4×10^6 Rhenium	76 Os 1×10^6 Osmium	77 Ir Iridium	78 Pt Platinum	79 Au 1×10^6 Gold	80 Hg 2×10^6 Mercury	81 Tl 1.5×10^6 Thallium	82 Pb 2×10^6 Lead	83 Bi 2×10^6 Bismuth	84 Po Polonium	85 At Astatine	
87 Fr Francium	88 Ra 1.5×10^6 Radium	89-103 Actinides	104 Rf Rutherfordium	105 Db Dubnium	106 Sg Seaborgium	107 Bh Bohrium	108 Hs Hassium	109 Mt Meitnerium	110 Ds Darmstadtium	111 Rg Roentgenium	112 Cn Copernicium	113 Uut Ununtrium	114 Fl Flerovium	115 Uup Ununpentium	116 Lv Livermorium	117 Uus Ununseptium	118 Uuo Ununoctium
Lanthanides		57 La 1×10^6 Lanthanum	58 Ce 1.2×10^6 Cerium	59 Pr 1×10^6 Praseodymium	60 Nd 2×10^6 Neodymium	61 Pm Promethium	62 Sm 4.5×10^6 Samarium	63 Eu 1.5×10^6 Europium	64 Gd 2×10^6 Gadolinium	65 Tb 1×10^6 Terbium	66 Dy 1.5×10^6 Dysprosium	67 Ho 2.2×10^6 Holmium	68 Er 1.7×10^6 Erbium	69 Tm 1.7×10^6 Thulium	70 Yb 1.2×10^6 Ytterbium	71 Lu 1.5×10^6 Lutetium	
Actinides		89 Ac Actinium	90 Th 1×10^6 Thorium	91 Pa 1×10^6 Protactinium	92 U 1.5×10^6 Uranium	93 Np Neptunium	94 Pu Plutonium	95 Am Americium	96 Cm Curium	97 Bk Berkelium	98 Cf Californium	99 Es Einsteinium	100 Fm Fermium	101 Md Mendelevium	102 No Nobelium	103 Lr Lawrencium	

Abundance Legend:

- $< 10^{-12}$
- $10^{-12} - 10^{-9}$
- $10^{-9} - 10^{-6}$
- $10^{-6} - 10^{-3}$
- $10^{-3} - 1$
- $1 - 10^3$
- $> 10^3$



	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1	H 1.00794																	He 4.002602
2	Li 6.941	Be 9.012182											B 10.811	C 12.0107	N 14.0067	O 15.9994	F 18.9984032	Ne 20.1797
3	Na 22.9897692	Mg 24.305											Al 26.9815386	Si 28.0855	P 30.973762	S 32.065	Cl 35.453	Ar 39.948
4	K 39.0983	Ca 40.078	Sc 44.9550	Ti 47.867	V 50.9415	Cr 51.9961	Mn 54.938045	Fe 55.845	Co 58.933195	Ni 58.6934	Cu 63.546	Zn 65.38	Ga 69.723	Ge 72.63	As 74.9216	Se 78.96	Br 79.904	Kr 83.798
5	Rb 85.4678	Sr 87.62	Y 88.90585	Zr 91.224	Nb 92.90638	Mo 95.94	Tc 98	Ru 101.07	Rh 102.9055	Pd 106.42	Ag 107.8682	Cd 112.411	In 114.818	Sn 118.71	Sb 121.76	Te 127.5	I 126.90447	Xe 131.293
6	Cs 132.905451	Ba 137.327		Hf 178.49	Ta 180.94788	W 183.84	Re 186.207	Os 190.23	Ir 192.22	Pt 195.084	Au 196.966569	Hg 200.59	Tl 204.3833	Pb 207.2	Bi 208.9804	Po 209	At 210	Rn 222
7	Fr 223	Ra 226		Rf 267	Db 268	Sg 271	Bh 272	Hs 270	Mt 278	Ds 281	Rg 280	Cn 285	Uut 284	Fl 289	Uup 288	Lv 293	Uus 294	Uuo 294
				La 138.90547	Ce 140.116	Pr 140.90765	Nd 144.242	Pm 145	Sm 150.36	Eu 151.964	Gd 157.25	Tb 158.92535	Dy 162.5	Ho 164.93032	Er 167.259	Tm 168.93421	Yb 173.054	Lu 174.9668
				Ac 227	Th 238.02891	Pa 231.03688	U 238.02891	Np 237	Pu 244	Am 243	Cm 247	Bk 247	Cf 251	Es 252	Fm 257	Md 258	No 259	Lr 262

Periodic Table of the Radioactive Elements

Half-life of Most Stable Isotope																	
Stable																	
$t_{1/2} > 10^8$ years																	
10^3 yrs $< t_{1/2} < 10^8$ yrs																	
1 yr $< t_{1/2} < 10^3$ yrs																	
1 day $< t_{1/2} < 10^1$ yrs																	
$t_{1/2} < 1$ day																	
unknown																	
1A	Atomic Number																8A
1	Symbol																2
H	# of isotopes																He
3	2A															10	
Li	Be															Ne	
11	12															18	
Na	Mg															Ar	
19	20	21	3B	4B	5B	6B	7B	8B		1B	2B	3A	4A	5A	6A	7A	8
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
55	56	57-71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ba	Lanthanides	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
87	88	89-103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118
Fr	Ra	Actinides	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Uut	Fl	Uup	Lv	Uus	Uuo
Lanthanides		57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	
		La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	
Actinides		89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	
		Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	

Temperature dependent Magnetic behaviour

$\text{NiCl}_2 \{ \text{P}(\text{C}_2\text{H}_5)_2(\text{C}_6\text{H}_5) \}_2$ exhibits temperature dependent magnetic behaviour (paramagnetic/diamagnetic). The coordination geometries of Ni^{2+} in the paramagnetic and diamagnetic states are respectively

- (a) tetrahedral and tetrahedral
 (b) square planar and square planar
 (c) tetrahedral and square planar
 (d) square planar and tetrahedral

Solution :



In the paramagnetic state, the complex will have unpaired 3d electrons in Ni^{2+} ion, whereas no such electrons (i.e. all the 3d electrons will be paired) are present in the diamagnetic state.

The electronic configuration of $_{28}\text{Ni}^{2+}$ is $\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\uparrow$ \square \square \square

If the two unpaired electrons remain present in the ion, it will undergo sp^3 hybridization to accommodate ligands and the geometry of the complex ion will be tetrahedral.

If the two unpaired electrons are coupled, the ion will be diamagnetic and it can undergo dsp^2 hybridization to accommodate ligands leading to the square planar geometry to the complex ion.

Therefore, the **choice (c)** is correct.




Reaction of white phosphorus with aqueous NaOH


The reaction of white phosphorus with aqueous NaOH gives phosphine along with another phosphorus containing compound. The reaction type, the oxidation states of phosphorus in phosphine and the other product, respectively, are

- (a) redox reaction, -3 and -5
(b) redox reaction, 3 and +5
(c) disproportionation reaction, -3 and +5
(c) disproportionation reaction, -3 and +3

Solution :



The salt NaH_2PO_2 undergoes the following changes on heating.



The reaction is disproportionation reaction and the oxidation states of phosphorus are -3 (in phosphine) and +5 in the second product.

Therefore, the **choice (c)** is correct.

Legend:

- Red circle: CCP: Cubic Close Packing, FCC: Face Centred Cubic
- Blue hexagon: hc (H4)
- Yellow square: BCC: Body Centred Cubic
- Green hexagon: HCP: Hexagonal Close Packing
- Yellow star: Other

Generally the stabilities of complexes decreases with increasing atomic number for the electropositive metals e.g. group IIA or hard Lewis acids, and increase with increasing atomic number for the noble metals (soft acids), following the general trend of ionization energy.

(a) For electropositive metals (hard acids) the order of stabilities of the Halide complexes $\text{F}^- > \text{Cl}^- > \text{Br}^- > \text{I}^-$ but for highly polarizing soft acid metal ions such as Hg^{+2} , we see the reverse order.

(b) The most electropositive metals (hardest acids) show a greater tendency of forming complexes, with hard ligands such as F^- or Oxygen containing ligands.

Electron affinities in the periodic table [hide]

Group →	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
↓ Period																		
1	H 79																	He *
2	Li 80	Be *											B 27	C 122	N *	O 141	F 328	Ne *
3	Na 53	Mg *											Al 42	Si 134	P 72	S 200	Cl 349	Ar *
4	K 48	Ca 2	Sc 18	Ti 8	V 51	Cr 65	Mn *	Fe 15	Co 64	Ni 112	Cu 119	Zn *	Ga 41	Ge 119	As 79	Se 195	Br 324	Kr *
5	Rb 47	Sr 5	Y 30	Zr 41	Nb 86	Mo 72	Tc *	Ru 101	Rh 110	Pd 54	Ag 126	Cd *	In 39	Sn 107	Sb 101	Te 190	I 295	Xe *
6	Cs 46	Ba 14	*	Hf 31	Ta 79	W 79	Re *	Os 104	Ir 150	Pt 205	Au 223	Hg *	Tl 36	Pb 35	Bi 91	Po *	At *	Rn *
7	Fr *	Ra *	**	Rf *	Db *	Sg *	Bh *	Hs *	Mt *	Ds *	Rg *	Cn *	Uut *	Fl *	Uup *	Lv *	Uus *	Uuo *
* Lanthanides			La 45	Ce 92	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm 99	Yb	Lu 33	
** Actinides			Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	

Legend
The number mentioned is **Electron affinity** in kJ/mol (rounded).
* Denotes elements that are expected to have electron affinities close to zero on quantum mechanical grounds

First Ionization Energies of Some Elements (kJ/mol)

period	Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1		H 1310																	He 2370
2		Li 520	Be 900											B 800	C 1090	N 1400	O 1310	F 1680	Ne 2080
3		Na 490	Mg 730											Al 580	Si 780	P 1060	S 1000	Cl 1250	Ar 1520
4		K 420	Ca 590	Sc 630	Ti 660	V 650	Cr 660	Mn 710	Fe 760	Co 760	Ni 730	Cu 740	Zn 910	Ga 580	Ge 780	As 960	Se 950	Br 1140	Kr 1350
5		Rb 400	Sr 550	Y 620	Zr 660	Nb 670	Mo 680	Tc 700	Ru 710	Rh 720	Pd 800	Ag 730	Cd 870	In 560	Sn 700	Sb 830	Te 870	I 1010	Xe 1170
6		Cs 380	Ba 500	La 540	Hf 700	Ta 760	W 770	Re 760	Os 840	Ir 890	Pt 870	Au 890	Hg 1000	Tl 590	Pb 710	Bi 800	Po 810	At ...	Rn 1030
7		Fr ...	Ra 510																

cyanide extraction process of silver from argentite ore


In the cyanide extraction process of silver from argentite ore, the oxidizing and reducing agents used are

- (a) O_2 and CO , respectively (b) O_2 and Zn dust, respectively
(c) HNO_3 and Zn, respectively (d) HNO_3 and CO , respectively

Solution :

The argentite ore contains Ag_2S . After crushing and concentration by froth flotation process, the ore is leached with 0.4 – 7% solution of sodium cyanide kept agitated with a current of air.

Silver passes into the solution as argentocyanide. $\text{Ag}_2\text{S} + 4\text{NaCN} \rightleftharpoons 2\text{Na}[\text{Ag}(\text{CN})_2] + \text{Na}_2\text{S}$
The air blown removes Na_2S as $\text{Na}_2\text{S}_2\text{O}_3$ and Na_2SO_4 causing the above reaction to proceed to completion. Silver is recovered by adding zinc into argentocyanide. $2\text{Ag}(\text{CN})_2^- + \text{Zn} \rightarrow [\text{Zn}(\text{CN})_4]^{2-} + 2\text{Ag}$
Thus, oxidizing agent is O_2 (from air) and the reducing agent is Zn.
Therefore, the **choice (b)** is correct.



Magnitude of electron affinity (kJ/mol),
s-, *p*-, and *d*-block elements

■ *s* block ■ *p* block ■ *d* block ■ *f* block

Common Name	Chemical Name
acetone	dimethyl ketone; 2-propanone (usually known as acetone)
acid potassium sulfate	potassium bisulfate
acid of sugar	oxalic acid
ackey	nitric acid
alcali volatil	ammonium hydroxide
alcohol, grain	ethyl alcohol
alcohol sulfuris	carbon disulfide
alcohol, wood	methyl alcohol
alum	aluminum potassium sulfate
alumina	aluminum oxide
antichlor	sodium thiosulfate
antimony black	antimony trisulfide
antimony bloom	antimony trioxide
antimony glance	antimony trisulfide
antimony red (vermillion)	antimony oxysulfide
aqua ammonia	aqueous solution of ammonium hydroxide
aqua fortis	nitric acid
aqua regia	nitrohydrochloric acid
aromatic spirit of ammonia	ammonia in alcohol
arsenic glass	arsenic trioxide
azurite	mineral form of basic copper carbonate
asbestos	magnesium silicate
aspirin	acetylsalicylic acid
baking soda	sodium bicarbonate
banana oil (artificial)	isoamyl acetate
barium white	barium sulfate
benzol	benzene
bicarbonate of soda	sodium hydrogen carbonate or sodium bicarbonate
bichloride of mercury	mercuric chloride
bichrome	potassium dichromate
bitter salt	magnesium sulfate
black ash	crude form of sodium carbonate

black copper oxide	cupric oxide
black lead	graphite (carbon)
blanc-fixe	barium sulfate
bleaching powder	chlorinated lime; calcium hypochlorite
blue copperas	copper sulfate (crystals)
blue lead	lead sulfate
blue salts	nickel sulfate
blue stone	copper sulfate (crystals)
blue vitriol	copper sulfate
bluestone	copper sulfate
bone ash	crude calcium phosphate
bone black	crude animal charcoal
boracic acid	boric acid
borax	sodium borate; sodium tetraborate
bremen blue	basic copper carbonate
brimstone	sulfur
burnt alum	anhydrous potassium aluminum sulfate
burnt lime	calcium oxide
burnt ochre	ferric oxide
burnt ore	ferric oxide
brine	aqueous sodium chloride solution
butter of antimony	antimony trichloride
butter of tin	anhydrous stannic chloride
butter of zinc	zinc chloride
calomel	mercury chloride; mercurous chloride
carbolic acid	phenol
carbonic acid gas	carbon dioxide
caustic lime	calcium hydroxide
caustic potash	potassium hydroxide
caustic soda	sodium hydroxide
chalk	calcium carbonate
Chile saltpeter	sodium nitrate
Chile nitre	sodium nitrate

Chinese red	basic lead chromate
Chinese white	zinc oxide
chloride of soda	sodium hypochlorite
chloride of lime	calcium hypochlorite
chrome alum	chromic potassium sulfate
chrome green	chromium oxide
chrome yellow	lead (VI) chromate
chromic acid	chromium trioxide
copperas	ferrous sulfate
corrosive sublimate	mercury (II) chloride
corundum (ruby, sapphire)	chiefly aluminum oxide
cream of tartar	potassium bitartrate
crocus powder	ferric oxide
crystal carbonate	sodium carbonate
dechlor	sodium thiophosphate
diamond	carbon crystal
emery powder	impure aluminum oxide
epsom salts	magnesium sulfate
ethanol	ethyl alcohol
farina	starch
ferro prussiate	potassium ferricyanide
ferrum	iron
flores martis	anhydride iron (III) chloride
fluorspar	natural calcium fluoride
fixed white	barium sulfate
flowers of sulfur	sulfur
'flowers of' any metal	oxide of the metal
formalin	aqueous formaldehyde solution
French chalk	natural magnesium silicate
French vergidris	basic copper acetate
galena	natural lead sulfide
Glauber's salt	sodium sulfate
green verditer	basic copper carbonate

green vitriol	ferrous sulfate crystals
gypsum	natural calcium sulfate
hard oil	boiled linseed oil
heavy spar	barium sulfate
hydrocyanic acid	hydrogen cyanide
hypo (photography)	sodium thiosulfate solution
Indian red	ferric oxide
Isinglass	agar-agar gelatin
jeweler's rouge	ferric oxide
killed spirits	zinc chloride
lampblack	crude form of carbon; charcoal
laughing gas	nitrous oxide
lead peroxide	lead dioxide
lead protoxide	lead oxide
lime	calcium oxide
lime, slaked	calcium hydroxide
limewater	aqueous solution of calcium hydroxide
liquor ammonia	ammonium hydroxide solution
litharge	lead monoxide
lunar caustic	silver nitrate
liver of sulfur	sulfurated potash
lye or soda lye	sodium hydroxide
magnesia	magnesium oxide
manganese black	manganese dioxide
marble	mainly calcium carbonate
mercury oxide, black	mercurous oxide
methanol	methyl alcohol
methylated spirits	methyl alcohol
milk of lime	calcium hydroxide
milk of magnesium	magnesium hydroxide
milk of sulfur	precipitated sulfur
"muriate" of a metal	chloride of the metal
muriatic acid	hydrochloric acid


natron	sodium carbonate
nitre	potassium nitrate
nordhausen acid	fuming sulfuric acid
oil of mars	deliquescent anhydrous iron (III) chloride
oil of vitriol	sulfuric acid
oil of wintergreen (artificial)	methyl salicylate
orthophosphoric acid	phosphoric acid
Paris blue	ferric ferrocyanide
Paris green	copper acetoarsenite
Paris white	powdered calcium carbonate
pear oil (artificial)	isoamyl acetate
pearl ash	potassium carbonate
permanent white	barium sulfate
plaster of Paris	calcium sulfate
plumbago	graphite
potash	potassium carbonate
potassa	potassium hydroxide
precipitated chalk	calcium carbonate
Prussic acid	hydrogen cyanide
pyro	tetrasodium pyrophosphate
quicklime	calcium oxide
quicksilver	mercury
red lead	lead tetraoxide
red liquor	aluminum acetate solution
red prussiate of potash	potassium ferrocyanide
red prussiate of soda	sodium ferrocyanide
Rochelle salt	potassium sodium tartrate
rock salt	sodium chloride
rouge, jeweler's	ferric oxide
rubbing alcohol	isopropyl alcohol
sal ammoniac	ammonium chloride
sal soda	sodium carbonate
salt, table	sodium chloride

salt of lemon	potassium binoxalate
salt of tartar	potassium carbonate
saltpeter	potassium nitrate
silica	silicon dioxide
slaked lime	calcium hydroxide
soda ash	sodium carbonate
soda nitre	sodium nitrate
soda lye	sodium hydroxide
soluble glass	sodium silicate
sour water	dilute sulfuric acid
spirit of hartshorn	ammonium hydroxide solution
spirit of salt	hydrochloric acid
spirit of wine	ethyl alcohol
spirits of nitrous ether	ethyl nitrate
sugar, table	sucrose
sugar of lead	lead acetate
sulfuric ether	ethyl ether
talc or talcum	magnesium silicate
tin crystals	stannous chloride
trona	natural sodium carbonate
unslaked lime	calcium oxide
Venetian red	ferric oxide
verdigris	basic copper acetate
Vienna lime	calcium carbonate
vinegar	impure dilute acetic acid
vitamin C	ascorbic acid
vitriol	sulfuric acid
washing soda	sodium carbonate
water glass	sodium silicate
white caustic	sodium hydroxide
white lead	basic lead carbonate
white vitriol	zinc sulfate crystals
yellow prussiate of potash	potassium ferrocyanide

yellow prussiate of soda sodium ferrocyanide

zinc vitriol zinc sulfate

zinc white zinc oxide



Colours of Compounds


Cr₂(SO₄)₃ - green



PbS - Black



[Fe(H₂O)₅NO]SO₄ - brown




(CH₃COO)₃Fe - blood red



NH₄Cl - white fumes





AgCl - curdy white precipitate



AgCl - curdy white precipitate

AgCl + NH₄OH → Ag(NH₃)₂Cl + H₂O



CrO₃ + HCl → CrO₂Cl₂ (deep red fumes) + H₂O



CrO₂Cl₂ (deep red fumes)



Na₂CrO₄ (yellow)




PbCrO₄ (yellow)





AgI, AgBr, AgCl





Ammonia test for silver halides. Test tubes and the results of using concentrated ammonia to test for the presence of three silver halides. Each one of the silver halides is next to a fuller test tube that shows the result of adding concentrated ammonia. Halides are compounds containing halogens such as here, from left to right silver chloride, silver bromide and silver iodide. Adding dilute ammonia dissolves the chloride, while concentrated ammonia is needed to dissolve the bromide, but it fails to dissolve the iodide. The silver halides are themselves the results of standard tests using silver nitrate to test for the halide ions.



BrCl - pale yellow



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



In the presence of Arsenic, we get yellow precipitate of $(\text{NH}_4)_3\text{AsO}_4 \cdot 12\text{MoO}_3$ (ammonium arsenomolybdate)



$\text{Mo}_3\text{O}_8 \cdot x\text{H}_2\text{O}$ - molybdenum blue



$\text{SiF}_4 + \text{H}_2\text{O} \rightarrow \text{H}_4\text{SiO}_4$ (silicic acid - white) + H_2SiO_4

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



$\text{Na}_2\text{S} + \text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}] \rightarrow \text{Na}_4[\text{Fe}(\text{CN})_5\text{NOS}]$ - violet colour



$\text{Na}_2\text{S} + \text{CdCO}_3 \rightarrow \text{CdS (yellow)} + \text{Na}_2\text{CO}_3$




$\text{Na}_3[\text{Co}(\text{NO}_2)_6] + \text{KCl} \rightarrow \text{K}_3[\text{Co}(\text{NO}_2)_6]$ (potassium cobaltnitrate - yellow ppt) + NaCl



AgCl , PbCl_2 , Hg_2Cl_2 - white

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



AgCl White

PbCl₂ is white



Hg₂Cl₂ is white



Hg₂Cl₂
white


H₂S, PbS, Bi₂S₃, CuS - black

H₂S is Black



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

PbS is Black



PbS Lead Sulfide is
Black

Bi₂S₃ is Black




CuS - black



Copper Sulfide
Powder

CdS, As₂S₃, SnS₂ - Yellow

CdS is Yellow



As₂S₃ is Yellow




SnS_2 is Yellow



Tin Sulfide SnS_2



Sb_2S_3 is orange




Sb_2S_3 is Orange

SnS is Brown



SnS is Brown is your teacher says so. SnS is Yellow if your teacher says so :-)

Cr(OH)₃ is green



Chromium (III) Oxide is Green




CoS, NiS - black

CoS is black



NiS is Black Mineral Millerite



Cobalt Oxide is Black



Cobalt Sulphate is deep red brown



Cobalt Nitrate is deep red brown



Cobalt Chloride is deep Brown red



Cobalt Acetate is deep Brown



Azulene - Deep Blue

Iodoform - CHI_3 Yellow

Iron Phenol complex - Violet - $[\text{Fe}(\text{OC}_6\text{H}_5)_6]^{-3}$

$6 \text{C}_6\text{H}_5\text{OH} + \text{FeCl}_3 \rightarrow [\text{Fe}(\text{OC}_6\text{H}_5)_6]^{-3} + 3 \text{H}^+ + \text{Cl}^-$

Ammonium Phosphomolybdate - Yellow - $(\text{NH}_4)_3\text{PO}_4 \cdot 12\text{MoO}_3$ (Yellow)

Prussian Blue - $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3 \cdot x\text{H}_2\text{O}$

$[\text{Fe}(\text{CN})_3\text{NOS}]^{-4}$ - Violet

$[\text{Fe}(\text{SCN})]^{2+}$ Blood Red



$[\text{Co}(\text{NH}_3)_5\text{Br}]^{+2} \text{SO}_4^{-2}$ - Violet - Pentaaminebromocobalt(III)sulphate

$[\text{Co}(\text{NH}_3)_5\text{SO}_4]^+ \text{Br}^- \rightarrow$ **Red**



Test for Halogen:

Halogens present in an organic compound forms sodium halide on fusion with sodium metal. Sodium halide extracted with water can be easily identified by adding silver nitrate solution after acidifying with dil. HNO_3 .



If chlorine is present, a white curdy precipitate soluble in ammonium hydroxide solution is formed.



If bromine is present, an yellowish white precipitate sparingly soluble in ammonium hydroxide solution is formed.



If iodine is present, an yellow precipitate insoluble in ammonium hydroxide solution is formed.




Test for Nitrogen:

The carbon and nitrogen present in the organic compound on fusion with sodium metal gives sodium cyanide (NaCN) soluble in water. This is converted in to sodium ferrocyanide by the addition of sufficient quantity of ferrous sulphate. Ferric ions generated during the process react with ferrocyanide to form prussian blue precipitate of ferric ferrocyanide.



Sodium ferrocyanide




Ferric ferrocyanide



Test for Sulphur:



If sulphur is present in the organic compound, sodium fusion will convert it into sodium sulphide. Sulphide ions are readily identified using sodium nitroprusside.




Sodium nitroprusside violet colour

Test for both Nitrogen and Sulphur:

If both nitrogen and sulphur are present in an organic compound, sodium fusion will convert it into sodium thiocyanate which then react with Fe^{3+} to form blood colour complex $[\text{Fe}(\text{SCN})]^{2+}$



Note: If the Lassaigne's extract containing excess of sodium metal, sodium cyanide and sulphides are formed instead of sodium thiocyanate.



Here in this type of cases, both sulphur and nitrogen are to be identified in separate tests.

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

$BaCO_3$, $SrCO_3$, $CaCO_3$, $Mg(NH_4)PO_4$ - white

$BaCO_3$ is white



$BaCO_3$ powder

$SrCO_3$ is white




$CaCO_3$ powder is white



$CaCO_3$ Calcium Carbonate Powder


CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

Mg(NH₄)PO₄ is white




Pentaamminechlorocobalt(III)Bromide

[Co(NH₃)₅Cl] Cl₂ Pentaamminechlorocobalt(III)chloride




Pentaamminesulphatocobalt(III)Bromide

Dark Yellow




trans- Pt(NH₃)₂Cl₂


Pale Yellow




cis- Pt(NH₃)₂Cl₂



Trans - [Co(NH₃)₄Cl₂]⁺





Cis - [Co(NH₃)₄Cl₂]⁺



Cis is Blue Violet


trans is green




Eta bonds - Also known as Hapticity - Hapticity is the coordination of a ligand to a metal center via an uninterrupted and contiguous series of atoms.

The hapticity of a ligand is described with the Greek letter η ('eta'). For example, η^2 describes a ligand that coordinates through 2 contiguous atoms. In general the η -notation only applies when multiple atoms are coordinated (otherwise the κ -notation is used). In addition, if the ligand coordinates through multiple atoms that are not contiguous then this is considered denticity (not hapticity), and the κ -notation is used once again. Lastly, bridging ligands are described with the μ ('mu') notation.

Ferrocene - bis(η^5 -cyclopentadienyl)iron



Uranocene - bis(η^8 -1,3,5,7-cyclooctatetraene)uranium




$W(CO)_3(PPr_i3)_2(\eta^2-H_2)$ - the first compound to be synthesized with a dihydrogen ligand (also known as Dihydrogen Complexes)
 $IrCl(CO)[P(C_6H_5)_3]_2(\eta^2-O_2)$ - the dioxygen derivative which forms reversibly upon oxygenation of Vaska's complex.

-


Chromium (III) chloride is purple



black ppt







$Bi(OH)_3$ Bismuth Hydroxide is white



Oxidation of Phenol



Test of Phenol

With Neutral $FeCl_3$, Phenol on reaction with neutral ferric chloride gives violet colour of ferric phenoxide






Liebermann's Nitroso Reaction



SbOCl Antimony oxychloride



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams


Antimony Trioxide is white



Antimony Pentoxide is gray




Antimony Trichloride is white




CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

Potassium antimonate white








Can Test

Treat a few drops of ceric ammonium nitrate with 3-4 ml of hot aqueous solution of the organic compound. A green or brown precipitate is formed.



Phenol gives blue colour with ammonia and sodium hypo chlorite.







CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



magnesium ammonium phosphate



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



Chromium oxide powder is green




Chromium Oxide
Powder

Bismuth oxide is Yellow



Bismuth Oxide
Powder

KI Potassium Iodide is white



CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams

Tin reacts with conc. HNO_3 forming metastannic acid (H_2SnO_3).

Tin is not attacked by organic acids and hence is used for tinning of utensils to resist corrosion. Tin foils are used for wrapping cigarettes, confectionary items and for making tooth-paste tubes.

SnO_2 is an amphoteric oxide.

Stannous chloride (SnCl_2) acts as a good reducing agent. It reduces HgCl_2 to first Hg_2Cl_2 and then to Hg . It also reduces FeCl_3 to FeCl_2 .

Stannic chloride (SnCl_4) is a liquid and fumes in air due to hydrolysis. It acts as a Lewis acid and dissolves in concentrated HCl forming H_2SnCl_6 .


$\text{SnCl}_4 \cdot 5\text{H}_2\text{O}$ is called butter of tin.

SnS dissolves in yellow ammonium sulphide.


Aluminum Oxide is white



Mercuric iodide HgI_2 : It is a yellow solid below 400K but changes to red solid above 400K.




It dissolves in excess of KI forming K_2HgI_4 ;



Alkaline solution of K_2HgI_4 is called Nessler's reagent.

Add NH_4OH excess and dimethylglyoxime to NiCl_2 then a rosy red ppt of nickel appears



A sample of $\text{Ni}(\text{dmgH})_2$


$\text{CuO} + \text{B}_2\text{O}_3$ (Glassy bead) \rightarrow $\text{Cu}(\text{BO}_2)_2$ (Copper metaborate - blue)



Compounds of zinc

Zinc oxide ZnO : Zincite (ZnO) is also called Philosopher's wool. It is a white powder, becomes **yellow** on heating and again white on cooling. It is amphoteric in nature. It is used as a white pigment under the name Zinc white or Chinese white.

Zinc Sulphate (white vitriol), $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$: It is a colourless transparent crystal highly soluble in water. It is used as an eye-lotion and for preparing double salts. On heating it loses its molecules of water as,



Verdigris - basic copper acetate



This pigment was used to the Middle Ages in paints. Until the nineteenth century this was the most vibrant shade of green available and it was used by artists widely. The color comes from copper acetate and you should know that it is highly toxic to humans.

Compounds of Mercury

(1) Mercuric oxide, HgO : It is obtained as a red solid by heating mercury in air or oxygen for a long time at 673 K



$2Hg + O_2 \rightarrow 2HgO$ (red) or by heating mercuric nitrate alone or in the presence of Hg


Heat



red




When NaOH is added to a solution of HgCl_2 , yellow precipitate of HgO are obtained.



(yellow)



Red and yellow forms of HgO differ only in their particle size. On heating to 673 K, yellow form changes to red form.



It is used in oil paints or as a mild antiseptic in ointments.

(2) Mercuric chloride, HgCl_2 : It is obtained by treating Hg with Cl_2 or by heating a mixture of NaCl and HgSO_4 in presence of small amount of MnO_2 (which oxidizes any Hg(I) salts formed during the reaction).

Heat



MnO_2







HgCl_2 is a white crystalline solid and is commonly known as corrosive sublimate. It is a covalent compound since it dissolves in organic solvents like ethanol and ether.

It is extremely poisonous and causes death. Its best antidote is white of an egg.

When treated with stannous chloride, it is first reduced to white ppt. of mercurous chloride and then to mercury (black).




white ppt.



grey


With ammonia it gives a white ppt. known as infusible white ppt.

$\text{HgCl}_2 + 2\text{NH}_3 \rightarrow \text{Hg}(\text{NH}_2)\text{Cl} + \text{NH}_4\text{Cl}$ A dilute solution of HgCl_2 is used as an antiseptic. (3) Mercuric iodide, HgI_2 : It is obtained when a required amount of KI solution is added to a solution of HgCl_2 . $\text{HgCl}_2 + 2\text{KI} \rightarrow \text{HgI}_2 + 2\text{KCl}$ (red) Below 400 K, HgI_2 is red but above 400 K, it turns yellow



An alkaline solution of $\text{K}_2[\text{HgI}_4]$ is called Nessler's reagent and is used to test NH_4^+ ions.

It gives a brown ppt. of $\text{NH}_2 - \text{Hg} - \text{O} - \text{Hg} - \text{I}$ (Iodide of Millon's base) with NH_4^+ ions.



It is used in ointments for treating skin infections.

(4) Mercurous chloride, Hg_2Cl_2 : It is obtained as under:



white ppt.

Heat in an iron retort



It is purified by sublimation.


Mercurous chloride is also called calomel. It is a white powder insoluble in H_2O . On heating, it decomposes to give HgCl_2 and Hg .



It dissolves in chlorine water forming mercuric chloride.



With ammonia, it turns black due to the formation of a mixture of finely divided black Hg and mercuric amino chloride.



(black)

It is used to prepare standard calomel electrode and as a purgative in medicine.

(5) Mercuric sulphide, HgS : The solubility product of HgS is lower than that of ZnS and hence it gets precipitated as black solid when H_2S is passed through an acidic solution of any mercury (II) salt.



It is insoluble in water and HCl but dissolves in aqua regia (1 part conc. HNO_3 + 3 parts conc. HCl)



Aqua regia Nitrosyl chloride Nascent chlorine




(Soluble)

On sublimation, its colour changes to red and hence it is used as a red pigment.

(6) Mercuric sulphate, HgSO_4 : It is obtained when HgS is treated with conc. H_2SO_4 .



It is a white solid which decomposes on heating to give mercurous sulphate.



It is used as a catalyst in the hydration of alkynes to give aldehydes or ketones. It is also used as a cosmetic under the name Vermillon and in ayurvedic medicine as makardhwaj.

(7) Amalgams : Mercury forms alloys commonly known as amalgams, with all metals except iron and platinum. Hence it is transported in iron containers.

(8) Alloy of transition metal : See in table discuss earlier in metallurgy.

XXXXXXXXXXXXXXXXXXXXXXX

The common Anions are divided into 3 groups for the purpose of identification:

Those which evolve gases with dilute hydrochloric acid :

Carbonate (CO_3^{2-})

bicarbonate(HCO_3^-)

Nitrite(NO_2^-)

Sulphide (S^{2-})

Sulphite(SO_3^{2-})

Thiosulphate($\text{S}_2\text{O}_3^{2-}$)

Those which do not react with dilute HCl, but which do evolve gases or volatile liquids with concentrated sulphuric acid:

Chloride (Cl^-)

Bromide (Br^-)

Iodide (I^-)

Nitrate (NO_3^-)

Those which do not react with either dilute hydrochloric acid or concentrated sulphuric acid:

Phosphate (PO_4^{3-})

Borate ($\text{B}_4\text{O}_7^{2-}$)

Sulphate (SO_4^{2-})

A Radical is an atom, or group of atoms that behaves like a single atom. Examples : H, K, Na, Cu, Pb, simple radicals ; NH_4^+ , SO_2 , NO_2 , compound

radicals.

A Basic Radical is a metal, or any radical behaving like a metal. Examples : Ag, Cu, Ca, Na, and NH_4 the last

so classified because it behaves like such simple radicals as Na, K, and Li.

An Acid Radical is a nonmetal, or any radical behaving like a nonmetal. Examples : Cl, Br, I, SO_2 , NO_2

An Acid is a substance containing H that may be replaced by a basic radical. When a nonmetal is combined with H only, as in HCl, HBr, H_2S , the compound is said to be a binary (i.e. composed of two elements) acid, or an hydracid. When the nonmetal is combined with H and O, the compound is called a ternary (i.e. composed of three or more elements) acid, or an oxyacid ; and the characteristic name of the acid is formed by adding the termination -ic to the name of the nonmetal to denote more oxygen, or the termination -ous to denote less. Examples : H_2SO_4 sulphuric, and H_2SO_3 sulphurous acid.

These are classified in six different groups in accordance with certain properties which permit a group precipitation by a single reagent. A reagent used for precipitation of a whole group is called a general reagent.

A reagent used as a test for a single substance is known as a special reagent.

Experiments showing Method of Classification

Use the following solutions : AgNO_3 , $\text{Cu}(\text{NO}_3)_2$, AsCl_3 , Fe_2Cl_6 , $\text{Co}(\text{NO}_3)_2$, $\text{Ba}(\text{NO}_3)_2$, NaCl.

Take about 2 c.c. of each solution, and add HCl.*

Points to be Observed Whether a precipitate is formed.

Whether the precipitate is soluble in an excess of the reagent.

* Always use dilute acids unless otherwise directed.

GROUPING THE BASES

The color of each precipitate.

Whether there is change in color of the precipitate.

Change in color of solutions.

Chemical equations.

Tabulate all results carefully in your notebook, according to the following scheme :

HCl was added to the seven solutions with the following results :

Typical of Gr. I $\text{AgNO}_3 + \text{HCl} = \text{AgCl} \downarrow + \text{HNO}_3$

Typical of Gr. II A $\text{Cu}(\text{NO}_3)_2 + \text{HCl} = \text{No change}$.

Typical of Gr. II B $\text{AsCl}_3 + \text{HCl} = \text{No change}$.

Typical of Gr. III $\text{FeCl}_2 + \text{HCl} = \text{No change}$.

Typical of Gr. IV $\text{Co}(\text{NO}_3)_2 + \text{HCl} = \text{No change}$.


Typical of Gr. V $\text{Ba}(\text{NO}_3)_2 + \text{HCl} = \text{No change}$.

Typical of Gr. VI $\text{NaCl} + \text{HCl} = \text{No change}$.

1) Anions which react with dilute hydrochloric acid

	Carbonates (CO_3^{2-})	Bicarbonates (HCO_3^-)	Nitrites (NO_2^-)
	All carbonates except those of alkali metals, and ammonium are very slightly or difficultly soluble in water. Accordingly reactions in solution are only carried out in case of the soluble salts.	All bicarbonates are water soluble.	Most nitrites are water soluble except that of silver.
Solid salt + dil. HCl	Effervescence and a colourless odourless gas is evolved, Carbon dioxide (CO_2), $\text{Na}_2\text{CO}_3 + 2\text{HCl} \rightarrow 2\text{NaCl} + \text{H}_2\text{O} + \text{CO}_2$	Effervescence and a colourless odourless gas is evolved, Carbon dioxide (CO_2), $\text{NaHCO}_3 + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O} + \text{CO}_2$	Pungent brown fumes are evolved consisting mainly of nitrogen dioxide (NO_2), which is produced by the combination of nitric oxide (NO) with the oxygen of the air. $\text{NaNO}_2 + \text{HCl} \rightarrow \text{NaCl} + \text{HNO}_2$ $3\text{HNO}_2 \rightarrow \text{HNO}_3 + \text{H}_2\text{O} + 2\text{NO}$ $2\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$

	Carbonate	Bicarbonate	Nitrites
To differentiate between CO_3^{2-} and HCO_3^-:			Confirmatory tests of nitrites:
1) Salt solution + magnesium sulphate solution	White precipitate of magnesium carbonate $\text{Na}_2\text{CO}_3 + \text{MgSO}_4 \rightarrow \text{MgCO}_3 \downarrow + \text{Na}_2\text{SO}_4$	No ppt. in the cold, as magnesium bicarbonate is soluble, but on heating, a white ppt. of magnesium carbonate is obtained: $2\text{NaHCO}_3 + \text{MgSO}_4 \rightarrow \text{Na}_2\text{SO}_4 + \text{Mg}(\text{HCO}_3)_2$ soluble in water $\text{Mg}(\text{HCO}_3)_2 \xrightarrow{\Delta} \text{MgCO}_3 \downarrow + \text{CO}_2 + \text{H}_2\text{O}$ ppt	1) The brown ring test: To a dilute solution of the nitrite, add two drops of freshly prepared ferrous sulphate solution (FeSO_4), and then add dil. H_2SO_4 very carefully drop by drop down the inside of the test tube: a brown ring forms due to the formation of $(\text{Fe,NO})\text{SO}_4$. Note: If the addition of dil. H_2SO_4 is not down the inside of the test tube, the solution becomes brown color (we can not notes the ring).
2) Salt solution + mercuric chloride solution	Reddish brown precipitate of mercuric carbonate $\text{Na}_2\text{CO}_3 + \text{HgCl}_2 \rightarrow \text{HgCO}_3 \downarrow + 2\text{NaCl}$	No ppt. in the cold, but on heating a reddish-brown ppt. of mercuric carbonate is obtained: $2\text{NaHCO}_3 + \text{HgCl}_2 \rightarrow 2\text{NaCl} + \text{Hg}(\text{HCO}_3)_2$ soluble in water $\text{Hg}(\text{HCO}_3)_2 \xrightarrow{\Delta} \text{HgCO}_3 \downarrow + \text{CO}_2 + \text{H}_2\text{O}$ ppt	2) solution of nitrite + potassium iodide solution (KI) + dilute sulfuric acid: a pale brown colour appears, due to the liberation of iodine. (The nitrite solution is an oxidizing agent.) $2\text{KNO}_2 + 2\text{KI} + 2\text{H}_2\text{SO}_4 \rightarrow 2\text{K}_2\text{SO}_4 + 2\text{H}_2\text{O} + \text{I}_2 + 2\text{NO}$ Iodide Iodine
			3) Solution of nitrite + acidified solution of potassium permanganate KMnO_4 (i.e. $\text{KMnO}_4 + \text{dil. H}_2\text{SO}_4$): the purple colour of the permanganate disappears. (the nitrite solution is an reducing agent.) $5\text{KNO}_2 + 2\text{KMnO}_4 + 3\text{H}_2\text{SO}_4 \rightarrow 5\text{KNO}_3 + \text{K}_2\text{SO}_4 + 2\text{MnSO}_4 + 3\text{H}_2\text{O}$



Is HgCO_3 reddish Brown ?



	Sulphites (SO_3^{2-})	Thiosulphates ($\text{S}_2\text{O}_3^{2-}$)	Sulphides (S^{2-})
	All sulphites are slightly soluble in water except the alkali metal sulphites and the ammonium salt.	Sodium thiosulphate is readily soluble in water, other thiosulphates are slightly soluble.	All sulphides, except those of alkali, alkali earth metals and the ammonium ion are slightly soluble in water.
Solid salt + dilute HCl	<p>Sulphur dioxide gas (SO_2), is evolved, which is recognizable by its pungent smell:</p> $\text{Na}_2\text{SO}_3 + 2\text{HCl} \rightarrow 2\text{NaCl} + \text{H}_2\text{O} + \text{SO}_2$ <p><u>Test for sulphur dioxide:</u> Dip a small piece of paper into an acidic solution of potassium dichromate, and hold it over the mouth of the test tube. The paper will turn green, because the sulphur dioxide reduces the dichromate to a chromic sulphate:</p> $3\text{SO}_2 + \text{K}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{SO}_4 \rightarrow \text{K}_2\text{SO}_4 + \text{Cr}_2(\text{SO}_4)_3 + \text{H}_2\text{O}$ <p>Orange colour Green colour</p>	<p>Colourless gas with pungent odour, which turns an acidified potassium dichromate paper green, is evolved (SO_2), and a yellow precipitate of sulphur (S) appears:</p> $\text{Na}_2\text{S}_2\text{O}_3 + 2\text{HCl} \rightarrow 2\text{NaCl} + \text{H}_2\text{O} + \text{SO}_2 + \text{S} \downarrow$ $3\text{SO}_2 + \text{K}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{SO}_4 \rightarrow \text{K}_2\text{SO}_4 + \text{Cr}_2(\text{SO}_4)_3 + \text{H}_2\text{O}$ <p>Orange colour Green colour</p>	<p>Colourless gas, hydrogen sulphide (H_2S), is evolved which is recognizable by its bad smell.</p> $\text{Na}_2\text{S} + 2\text{HCl} \rightarrow 2\text{NaCl} + \text{H}_2\text{S}$ <p><u>Test for hydrogen sulphide:</u> Dip a piece of paper in a solution of lead acetate, and hold it over the mouth of the test tube in which the H_2S is being generated. The paper will turn black due to the formation of lead sulphide.</p> $\text{H}_2\text{S} + \text{Pb}(\text{CH}_3\text{COO})_2 \rightarrow \text{PbS} + 2\text{CH}_3\text{COOH}$ <p>Black</p>



K_2SO_4 is white



$\text{Cr}_2(\text{SO}_4)_3$ is green


	Sulphites	Thiosulphates	Sulphides
Confirmatory tests: 1) Salt solution + silver nitrate solution (AgNO ₃)	A white precipitate forms (silver sulphite Ag ₂ SO ₃), which dissolves in excess of sulphite due to the formation of soluble complex. $\text{Na}_2\text{SO}_3 + 2 \text{AgNO}_3 \rightarrow \text{Ag}_2\text{SO}_3 \downarrow + 2 \text{NaNO}_3$ ppt $\text{Ag}_2\text{SO}_3 + \text{Na}_2\text{SO}_3 \rightarrow 2 \text{Na}[\text{AgSO}_3]$ Soluble complex	A white precipitate forms (silver thiosulphate Ag ₂ S ₂ O ₃), which is soluble in excess of the thiosulphate due to the formation of complex, which is unstable (it changes to yellow, brown and finally to black Ag ₂ S). $\text{Na}_2\text{S}_2\text{O}_3 + 2 \text{AgNO}_3 \rightarrow \text{Ag}_2\text{S}_2\text{O}_3 \downarrow + 2 \text{NaNO}_3$ White ppt	A black precipitate of silver sulphide (Ag ₂ S) is formed. $\text{Na}_2\text{S} + 2\text{AgNO}_3 \rightarrow \text{Ag}_2\text{S} \downarrow + 2\text{NaNO}_3$ ppt
2) Salt solution + lead acetate solution Pb(CH ₃ COO) ₂	A white ppt. of lead sulphite (PbSO ₃) is formed. $\text{Na}_2\text{SO}_3 + \text{Pb}(\text{CH}_3\text{COO})_2 \rightarrow \text{PbSO}_3 \downarrow + 2 \text{CH}_3\text{COONa}$ ppt	A white ppt. of lead thiosulphate (PbS ₂ O ₃) is formed, the ppt. turns black (PbS) on boiling. $\text{Na}_2\text{S}_2\text{O}_3 + \text{Pb}(\text{CH}_3\text{COO})_2 \rightarrow \text{PbS}_2\text{O}_3 \downarrow + 2 \text{CH}_3\text{COONa}$ White ppt $\text{PbS}_2\text{O}_3 + \text{H}_2\text{O} \rightarrow \text{PbS} \downarrow + \text{H}_2\text{SO}_4$ Black ppt	A black ppt. of lead sulphide (PbS) is formed. $\text{Na}_2\text{S} + \text{Pb}(\text{CH}_3\text{COO})_2 \rightarrow \text{PbS} \downarrow + 2 \text{CH}_3\text{COONa}$ Black ppt
	* Salt solution + drops of acidified solution of KMnO ₄ : The purple colour of the permanganate will disappear. The sulphite reduces the permanganate ion to (Mn ²⁺) which is almost colourless. $5\text{Na}_2\text{SO}_3 + 2\text{KMnO}_4 + 3\text{H}_2\text{SO}_4 \rightarrow 5\text{Na}_2\text{SO}_4 + \text{K}_2\text{SO}_4 + 2\text{MnSO}_4 + 3\text{H}_2\text{O}$		* Salt solution + Sodium nitroprusside solution: A violet colour is formed.

	Sulphites	Thiosulphates
	Salt solution + 2 drops of dil. H₂SO₄ + Iodine solution (I₂)	
	The brown colour of the iodine solution disappears. $\text{Na}_2\text{SO}_3 + \text{I}_2 + \text{H}_2\text{O} \rightarrow \text{Na}_2\text{SO}_4 + 2\text{HI}$	The brown colour of the iodine solution disappears, because the iodine is reduced to the iodide ion, which is colourless. At the same time, the thiosulphate is oxidized to tetrathionate: $2\text{Na}_2\text{S}_2\text{O}_3 + \text{I}_2 \rightarrow \text{Na}_2\text{S}_4\text{O}_6 + 2 \text{NaI}$


2) Anions which react with concentrated sulphuric acid

Note: Do these tests in fuming cuper because the gases which are given off in these tests are extremely irritating, and can cause damage to the sensitive mucous membranes of nose and throat.


	Chlorides (Cl ⁻)	Bromides (Br ⁻)	Iodides (I ⁻)	Nitrates (NO ₃ ⁻)
	All chlorides are water soluble except the chlorides of silver, mercurous and cuprous.	Bromides resemble chlorides in their solubility.	Iodides resemble chlorides and bromides in their solubility. However, bismuth iodide is insoluble.	All nitrates are soluble in water except some basic nitrates.
Solid salt + conc. H₂SO₄:	Effervescence with evolution of colourless gas, Hydrogen chloride (HCl). $2\text{NaCl} + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + 2\text{HCl}$ Test for HCl gas: Dip a clean glass rod into a bottle of conc. ammonia solution, and hold it close to the mouth of the test tube in which the HCl is being produced. Dense white fumes of ammonium chloride will be formed: $\text{HCl} + \text{NH}_3 \rightarrow \text{NH}_4\text{Cl}$	Reddish fumes evolve and the solution turns orange due to liberation of bromine (Br ₂). $2\text{NaBr} + \text{H}_2\text{SO}_4 \rightarrow 2\text{HBr} + \text{Na}_2\text{SO}_4$ $2\text{HBr} + \text{H}_2\text{SO}_4 \rightarrow 2\text{H}_2\text{O} + \text{SO}_2 + \text{Br}_2$	Violet fumes are evolved, (Iodine gas I ₂) and a brown or black precipitate is formed in the test tube. $2\text{KI} + 2\text{H}_2\text{SO}_4 \rightarrow \text{K}_2\text{SO}_4 + 2\text{H}_2\text{O} + \text{SO}_2 + \text{I}_2$	Yes If a small piece of copper metal and drops of water are added, dense brown fumes of nitrogen dioxide (NO ₂) will be given off. If no results, heat carefully. $2\text{KNO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{K}_2\text{SO}_4 + 2\text{HNO}_3$ $4\text{HNO}_3 + \text{Cu} \rightarrow \text{Cu}(\text{NO}_3)_2 + 2\text{H}_2\text{O} + 2\text{NO}_2$

	Chlorides	Bromides	Iodides	Nitrates
Confirmatory tests: 1) Salt solution + silver nitrate (AgNO ₃)	A dense white ppt. of silver chloride (AgCl) slowly turns a violet colour when exposed to bright sunlight. $\text{NaCl} + \text{AgNO}_3 \rightarrow \text{AgCl} \downarrow + \text{NaNO}_3$ ppt	A yellowish white precipitate of silver bromide (AgBr) is formed. $\text{NaBr} + \text{AgNO}_3 \rightarrow \text{AgBr} \downarrow + \text{NaNO}_3$ ppt	A yellow precipitate of silver iodide (AgI) is formed. $\text{KI} + \text{AgNO}_3 \rightarrow \text{AgI} \downarrow + \text{KNO}_3$ ppt	Yes
2) Salt solution + lead acetate Pb(CH ₃ COO) ₂	A white precipitate of lead chloride (PbCl ₂) is formed which is soluble in hot water, and reprecipitates on cooling. $2\text{NaCl} + \text{Pb}(\text{CH}_3\text{COO})_2 \rightarrow \text{PbCl}_2 \downarrow + 2\text{CH}_3\text{COONa}$	A white precipitate of lead bromide (PbBr ₂) appears which is soluble in boiling water and reprecipitates on cooling. $2\text{NaBr} + \text{Pb}(\text{CH}_3\text{COO})_2 \rightarrow \text{PbBr}_2 \downarrow + 2\text{CH}_3\text{COONa}$	A yellow precipitate of lead iodide (PbI ₂) is formed which dissolves in boiling water and recrystallises on cooling. $2\text{KI} + \text{Pb}(\text{CH}_3\text{COO})_2 \rightarrow \text{PbI}_2 \downarrow + 2\text{CH}_3\text{COOK}$	
			* Salt solution + Mercuric chloride (HgCl₂): A reddish precipitate of mercuric iodide is formed which dissolves in excess of KI. $\text{HgCl}_2 + 2\text{KI} \rightarrow \text{HgI}_2 \downarrow + 2\text{KCl}$ ppt $\text{HgI}_2 + 2\text{KI} \rightarrow \text{K}_2[\text{HgI}_4]$ soluble	* Brown ring test: The nitrate solution is mixed with freshly prepared FeSO ₄ solution, then conc. H ₂ SO ₄ is added and allowed to flow cautiously on the side of the test tube. A brown ring (Fe(NO)SO ₄) is formed at the interface of the two layers. The brown ring disappears on shaking the solution.
			* Salt solution + copper sulphate (CuSO₄): A brown ppt. of cuprous iodide CuI and iodine I ₂ . $4\text{KI} + 2\text{CuSO}_4 \rightarrow 2\text{CuI} \downarrow + 2\text{K}_2\text{SO}_4 + \text{I}_2$	


AgCl is white

CuI Cuprous iodide




Brown Ring Test



$$2\text{NaNO}_3 + \text{H}_2\text{SO}_4 \longrightarrow 2\text{NaHSO}_4 + \text{HNO}_3;$$

$$2\text{HNO}_3 + 6\text{FeSO}_4 + 3\text{H}_2\text{SO}_4 \longrightarrow 3\text{Fe}_2(\text{SO}_4)_3 + 2\text{NO} + 4\text{H}_2\text{O}$$

$$[\text{Fe}(\text{H}_2\text{O})_6]\text{SO}_4 + \text{NO} \longrightarrow [\text{Fe}(\text{H}_2\text{O})_5\text{NO}]^{2+} \text{SO}_4^{2-} + \text{H}_2\text{O}$$



3) Anions which do not react with acids:


	Phosphates (H_2PO_4^- , HPO_4^{2-} , PO_4^{3-})	Borates (BO_2^- , BO_3^{--} , $\text{B}_4\text{O}_7^{--}$)	Sulphates (SO_4^{2-})
	Most phosphates are insoluble in water except those of ammonium and alkali metals.	Ammonium and alkali metal borates are water soluble while other borates are slightly soluble in water.	All sulphates are soluble in water except those of some divalent metals e.g. calcium, strontium, barium and lead.
Salt solution + Barium chloride solution (BaCl_2)	A white precipitate of barium phosphate (BaHPO_4) is produced, soluble in dilute acids e.g. HNO_3 or HCl and insoluble in excess of barium chloride. $\text{Na}_2\text{HPO}_4 + \text{BaCl}_2 \rightarrow \text{BaHPO}_4\downarrow + 2\text{NaCl}$	A white ppt. of barium borate from concentrated solutions $\text{Ba}(\text{BO}_2)_2$ is produced, soluble in dilute acids and in excess of barium chloride. $\text{Na}_2\text{B}_4\text{O}_7 + 3\text{H}_2\text{O} + \text{BaCl}_2 \rightarrow \text{Ba}(\text{BO}_2)_2 + 2\text{H}_3\text{BO}_3 + 2\text{NaCl}$	A white precipitate of barium sulphate (BaSO_4) is formed which is insoluble in dilute acids and in excess of barium chloride. $\text{Na}_2\text{SO}_4 + \text{BaCl}_2 \rightarrow \text{BaSO}_4\downarrow + 2\text{NaCl}$
Confirmatory tests: 1) Salt solution + silver nitrate (AgNO_3)	A yellow precipitate of silver phosphate (Ag_3PO_4) is formed, which is readily soluble in dil. HNO_3 and ammonia.	A white ppt. of silver borate (AgBO_2) is formed, from concentrated solution, which give brown ppt. after boiling. (also a brown ppt. is formed with diluted solution) $\text{Na}_2\text{B}_4\text{O}_7 + 3\text{H}_2\text{O} + 2\text{AgNO}_3 \rightarrow 2\text{AgBO}_2 + 2\text{H}_3\text{BO}_3 + 2\text{NaNO}_3$ $2\text{AgBO}_2 + 3\text{H}_2\text{O} \rightarrow \text{Ag}_2\text{O} + 2\text{H}_3\text{BO}_3$	A white ppt. of silver sulphate (Ag_2SO_4) is formed with concentrated solution. $\text{Na}_2\text{SO}_4 + 2\text{AgNO}_3 \rightarrow \text{Ag}_2\text{SO}_4\downarrow + 2\text{NaNO}_3$



BaSO₄ is white



<p>For phosphates:</p> <p><u>0.5 ml of salt solution + 4 ml of ammonium molybdate solution + 0.5 ml of conc. HNO₃:</u></p> <p>Form a canary yellow precipitate of ammonium phosphomolybdate (NH₄)₃PO₄ · 12MoO₃ · H₂O in the cold or by gentle warming. (note the volume needed in this test).</p>		<p>For sulphates:</p> <p><u>Salt solution + lead acetate:</u></p> <p>A white precipitate of lead sulphate is formed, which is readily soluble in hot concentrated ammonium acetate or conc. H₂SO₄</p> <p>$K_2SO_4 + Pb(CH_3COO)_2 \rightarrow PbSO_4 \downarrow + 2CH_3COOK$</p> <p>$PbSO_4 + H_2SO_4 \rightarrow Pb(HSO_4)_2$</p>
--	--	---



Scheme for the identification of acid radicals

1) Solid salt + dil. HCl:

Observation	Effervescence and a colourless odourless gas is evolved.	Pungent brown fumes are evolved.	Colourless gas with Pungent odour, which turns an acidified dichromate paper green, is evolved.	Colourless gas with pungent odour, which turns an acidified potassium dichromate paper green, is evolved, and a yellow precipitate of sulphur appears.	Colourless gas is evolved which is characterized by its bad odour and turning a lead acetate paper black.	-ve The salt does not react with dil. HCl.
Results	The gas is CO_2 . The anion is carbonate or bicarbonate.	The gas is NO_2 . The anion is Nitrite.	The gas is SO_2 and the green colour is $\text{Cr}_2(\text{SO}_4)_3$. The anion is sulphite.	The gas is SO_2 , the ppt. is S and the green colour is $\text{Cr}_2(\text{SO}_4)_3$. The anion is thiosulphate.	The gas is H_2S , and the black colour is PbS. The anion is sulphide.	The anion is not from group 1. <u>(Go to step 2).</u>

To differentiate between carbonate and bicarbonate:

<p>1) Salt solution + magnesium sulphate solution: For carbonate: White precipitate of magnesium carbonate (MgCO_3). For bicarbonate: No ppt. in the cold, as magnesium bicarbonate is soluble, but on heating, a white ppt of magnesium carbonate is obtained.</p>	<p>2) Salt solution + mercuric chloride solution: For carbonate: Reddish brown precipitate of mercuric carbonate (HgCO_3). For bicarbonate: No ppt. in the cold, but on heating a reddish-brown ppt. of mercuric carbonate is obtained.</p>
---	---

2) Solid salt + conc. H_2SO_4 :

Observation	Effervescence with the evolution of a colourless gas which forms white fumes with NH_4OH .	Reddish fumes evolve and the solution turns orange.	Violet fumes are evolved, and a brown or black precipitate is formed in the test tube.	-ve If a small piece of copper metal and drops of water are added, dense brown fumes will be given off. If no results, heat carefully.	-ve The salt does not react with conc. H_2SO_4 .
Results	The gas is HCl and the white fumes are NH_4Cl . The anion is chloride.	The gas is Br_2 . The anion is Bromide.	The gas is I_2 . The anion is iodide.	The gas is NO_2 . The anion is Nitrate.	The anion is not from group 2. <u>(Go to step 3).</u>

3) Salt solution + BaCl_2 :

Observation	A white precipitate is produced, soluble in dilute acids e.g. HNO_3 or HCl and insoluble in excess of barium chloride.	A white ppt. from concentrated solutions is produced, soluble in dilute acids and in excess of barium chloride.	A white precipitate is formed which is insoluble in dilute acids and in excess of barium chloride.
Results	The white precipitate is barium phosphate (BaHPO_4). The anion is phosphate.	The white ppt. is barium borate $\text{Ba}(\text{BO}_2)$. The anion is Borate.	A white precipitate is barium sulphate (BaSO_4). The anion is sulphate.

<u>To differentiate between phosphate, borate and sulphate:</u>			
<u>Salt solution + AgNO₃ (silver nitrate):</u>			
	phosphate	borate	Sulphate
Observation	A yellow precipitate is formed, which is readily soluble in dil. HNO ₃ and ammonia.	A white ppt. is formed from concentrated solution, which give brown ppt. after boiling. (also, a brown ppt. is formed with diluted solution.)	A white ppt. is formed, with concentrated solution.
Results	The yellow ppt. is silver phosphate (Ag ₃ PO ₄). The anion is phosphate.	The white ppt. is silver borate (AgBO ₂). The anion is borate.	The white ppt. is silver sulphate (Ag ₂ SO ₄). The anion is sulphate.

Oxidation States and Colors

Metal	+1	+2	+3	+4	+5	+6	+7
Sc			Colorless				
Ti			Violet	Colorless			
V		Violet	Green	blue	Yellow		
Cr		Blue	Green			Yellow (CrO_4^{2-}) orange ($\text{Cr}_2\text{O}_7^{2-}$)	
Mn		Pale pink		Brown		Dark green (MnO_4^{2-})	Intense purple (MnO_4^-)
Fe		Pale green	Yellow or brown				
Co		Pink	Orange/y ellow				
Ni		Green					
Cu		Blue					
Zn		colorless					

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams



(In 2016 Celebrating 27 years of Excellence in Teaching)

Good Luck to you for your Preparations, References, and Exams

All Other Books written by me can be downloaded from

<https://zookeepersblog.wordpress.com/free-pdf-e-book-download-for-iit-jee-cbse-isc-cet-physics-chemistry-maths-from-professor-subhashish-skmclasses/>

Professor [Subhashish Chattopadhyay](#)

Learn more at <http://skmclasses.weebly.com/iit-jee-home-tuitions-bangalore.html>

Twitter - <https://twitter.com/ZookeeperPhy>

Facebook - <https://www.facebook.com/IIT.JEE.by.Prof.Subhashish/>

Blog - <http://skmclasses.kinja.com>

Blog - <http://skmclasses.blog.com>

CBSE Standard 12 Chemistry Survival Guide - Reduction Methods by Prof. Subhashish Chattopadhyay
SKMClasses Bangalore Useful for IIT-JEE, I.Sc. PU-II, Boards, IGCSE IB AP-Chemistry and other exams